



US 20020045607A1

(19) **United States**(12) **Patent Application Publication** (10) Pub. No.: **US 2002/0045607 A1**
Beachy et al. (43) Pub. Date: **Apr. 18, 2002**(54) **CHOLESTEROL AND HEDGEHOG
SIGNALING**sional of provisional application No. 60/074,714,
filed on Feb. 13, 1998.(75) Inventors: **Philip A. Beachy**, Baltimore, MD (US);
Jeffrey A. Porter, Belmont, MA (US);
Michael K. Cooper, Baltimore, MD
(US)**Publication Classification**(51) Int. Cl.⁷ **A61K 31/58**
(52) U.S. Cl. **514/176**

Correspondence Address:

LISA A. HAILE, Ph.D.**GRAY CARY WARE & FREIDENRICH LLP****4365 Executive Drive, Suite 1100****San Diego, CA 92121-2133 (US)**(57) **ABSTRACT**(73) Assignee: **The Johns Hopkins University School
of Medicine**(21) Appl. No.: **09/954,727**(22) Filed: **Sep. 11, 2001****Related U.S. Application Data**(63) Continuation of application No. 09/250,785, filed on
Feb. 12, 1999, now patented, which is a non-provi-

The present invention sterol-modified hedgehog polypeptides and functional fragments thereof. Methods of identifying compositions which affect hedgehog activity based on inhibition of cholesterol modification of hedgehog protein are described. In one aspect of the invention, the method provides a means for affecting cholesterol biosynthesis or transport in a cell comprising contacting a cell with an effective amount of a compound that affects hedgehog, thereby affecting cholesterol biosynthesis or transport. The effect may be inhibition or stimulation of cholesterol biosynthesis or transport.



US 20020006931A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2002/0006931 A1**
BEACHY et al. (43) **Pub. Date: Jan. 17, 2002**

(54) **INHIBITORS OF HEDGEHOG SIGNALING
PATHWAYS, COMPOSITIONS AND USES
RELATED THERETO**

(22) Filed: **Jun. 4, 1998**

Related U.S. Application Data

(76) Inventors: **PHILIP A. BEACHY, BALTIMORE,
MD (US); MICHAEL K. COOPER,
BALTIMORE, MD (US); JEFFREY A.
PORTER, CAMBRIDGE, MA (US)**

(63) Non-provisional of provisional application No.
60/081,186, filed on Apr. 9, 1998.

Publication Classification

Correspondence Address:

**ROPES & GRAY
ONE INTERNATIONAL PLACE
BOSTON, MA 02110-2624 (US)**

(51) **Int. Cl.⁷ A61K 31/56; A61K 31/44**

(52) **U.S. Cl. 514/278**

(57) **ABSTRACT**

(*) Notice: This is a publication of a continued prosecution application (CPA) filed under 37 CFR 1.53(d).

The present invention makes available assays and reagents inhibiting paracrine and/or autocrine signals produced by a hedgehog protein comprising contacting a cell sensitive to the hedgehog protein with a steroidal alkaloid, or other small molecule, in a sufficient amount to reduce the sensitivity of the cell to the hedgehog protein.

(21) Appl. No.: **09/090,622**



US 20020068706A1

(19) **United States**

(12) **Patent Application Publication**
GYURIS et al.

(10) **Pub. No.: US 2002/0068706 A1**

(43) **Pub. Date: Jun. 6, 2002**

(54) **INHIBITORS OF CELL-CYCLE
PROGRESSION AND USES RELATED
THERE TO**

(22) Filed: **Jul. 29, 1997**

Related U.S. Application Data

(76) Inventors: **JENO GYURIS, WINCHESTER, MA
(US); LOU LAMPHERE, BOSTON,
MA (US); DAVID H. BEACH,
HUNTINGTON BAY, NY (US)**

(63) Continuation-in-part of application No. 08/589,981,
filed on Jan. 23, 1996, now Pat. No. 5,672,508.

Publication Classification

Correspondence Address:

**FOLEY,HOAG & ELIOT, LLP
PATENT GROUP
ONE POST OFFICE SQUARE, LLP
BOSTON, MA 02109 (US)**

(51) **Int. Cl.⁷ A61K 31/70; C12N 15/63;
C07H 21/04**

(52) **U.S. Cl. 514/44; 435/455; 536/23.4;
536/23.72; 536/24.1**

(*) Notice: This is a publication of a continued prosecution application (CPA) filed under 37 CFR 1.53(d).

(57) **ABSTRACT**

The present invention pertains to novel inhibitors of cyclin-dependent kinases (CDKs), particularly CDK/cyclin complexes, which inhibitors can be used to control proliferation and/or differentiation of cells in which the inhibitors are introduced.

(21) Appl. No.: **08/902,572**

INHIBITORS OF CELL-CYCLE PROGRESSION AND USES RELATED THERETO

RELATED APPLICATIONS

[0001] This application is a continuation-in-part of U.S. Ser. No. 08/589,981, entitled *Inhibitors of Cell-Cycle Progression, and Uses Related Thereto*, filed on Jan. 23, 1996.

BACKGROUND OF THE INVENTION

[0002] The cell division cycle is one of the most fundamental processes in biology which, in multicellular organisms, ensures the controlled generation of cells with specialized functions. Under normal growth conditions, cell proliferation is tightly regulated in response to diverse intra- and extracellular signals. This is achieved by a complex network of proto-oncogenes and tumor-suppressor genes that are components of various signal transduction pathways. Activation of a proto-oncogene(s) and/or a loss of a tumor suppressor gene(s) can lead to the unregulated activity of the cell cycle machinery. This, in turn, will lead to unregulated cell proliferation and to the accumulation of genetic errors which ultimately will result in the development of cancer (Pardee, *Science* 246:603-608, 1989).

[0003] In the eukaryotic cell cycle a key role is played by the cyclin-dependent kinases (CDKs). Cdk complexes are formed via the association of a regulatory cyclin subunit and a catalytic kinase subunit. In mammalian cells, the combination of the kinase subunits (such as cdc2, CDK2, CDK4 or CDK6) with a variety of cyclin subunits (such as cyclin A, B1, B2, D1, D2, D3 or E) results in the assembly of functionally distinct kinase complexes. The coordinated activation of these complexes drives the cells through the cell cycle and ensures the fidelity of the process (Draetta, *Trends Biochem. Sci.* 15:378-382, 1990; Sherr, *Cell* 73:1059-1065, 1993). Each step in the cell cycle is regulated by a distinct and specific cyclin-dependent kinase. For example, complexes of Cdk4 and D-type cyclins govern the early G1 phase of the cell cycle, while the activity of the CDK2/cyclin E complex is rate limiting for the G1 to S-phase transition. The CDK2/cyclin A kinase is required for the progression through S-phase and the cdc2/cyclin B complex controls the entry into M-phase (Sherr, *Cell* 73:1059-1065, 1993).

[0004] The CDK complex activity is regulated by mechanisms such as stimulatory or inhibitory phosphorylations as well as the synthesis and degradation of the kinase and cyclin subunit themselves. Recently, a link has been established between the regulation of the activity of cyclin-dependent kinases and cancer by the discovery of a group of CDK inhibitors including the p27^{Kip1}, p21^{Waf1/Cip1} and p16^{Ink4/MTS1} proteins. The activity of p21^{Waf1/Cip1} is regulated transcriptionally by DNA damage through the induction of p53, senescence and quiescence (Harper et al., *Cell* 75:805-816, 1993). The inhibitory activity of p27^{Kip1} is induced by the negative growth factor TGF- β and by contact inhibition (Polyak et al., *Cell* 78:66-69, 1994). These proteins, when bound to CDK complexes, inhibit their kinase activity, thereby inhibiting progression through the cell cycle. Although their precise mechanism of action is unknown, it is thought that binding of these inhibitors to the CDK/cyclin complex prevents its activation. Alternatively, these inhibitors may interfere with the interaction of the enzyme with its substrates or its cofactors.

[0005] While p21^{Waf1/Cip1} and p27^{Kip1} inhibit all the CDK/cyclin complexes tested, p16^{Ink4/MTS1}, p15, p18 and p19 block exclusively the activity of the CDK4/cyclin D and CDK6/cyclin D complexes in the early G1 phase (Serrano et al., *Nature* 366:704-707, 1993), by either preventing the interaction of Cdk4 and Cyclin D1, or indirectly preventing catalysis. As mentioned above, the p21^{Waf1/Cip1} is positively regulated by the tumor suppressor p53 which is mutated in approx. 50% of all human cancers. p21^{Waf1/Cip1} may mediate the tumor suppressor activity of p53 at the level of cyclin-dependent kinase activity. p16^{Ink4/MTS1} is the product of a tumor suppressor gene localized to the 9p21 locus, which is frequently mutated in human cancer cells.

[0006] Of all the various kinases, the CDK4/cyclin D complexes are known to play an important role in regulating cell cycle progression in early G1. These complexes function as integrators of various growth factor-induced extracellular signals and as a link between the different signal transduction pathways and other cyclin-dependent kinases. The expression of the cyclin D1 positive regulatory subunit, is deregulated by gene translocations, retroviral insertions and amplifications in parathyroid adenomas, lymphomas, esophageal and breast carcinomas. The targeted overexpression of cyclin D1 in the mammary epithelium of transgenic mice induces mammary adenomas and adenocarcinomas. This confirms that cyclin D1, when overexpressed, acts as an oncogene (Wang et al., *Nature* 369:669-671, 1994). These data supports the idea that the lack of functional p16^{Ink4/MTS1} or the overexpression of cyclin D1 leads to the deregulation of CDK4/cyclin D1 kinase activity and thereby contribute to uncontrolled cell proliferation.

[0007] The prominent role of CDK/cyclin kinase complexes, in particular, CDK4/cyclin D kinase complexes, in the induction of cell proliferation and their deregulation in tumors, makes them ideal targets for developing highly specific anti-proliferative agents.

SUMMARY OF THE INVENTION

[0008] In one aspect, the present invention relates to a nucleic acid comprising a nucleotide sequence encoding a chimeric polypeptide having at least two CDK-binding motifs derived from different proteins which bind to cyclin dependent kinases (CDKs). The chimeric polypeptide binds to CDKs and inhibits cell-cycle progression.

[0009] The chimeric polypeptide can be a fusion protein, or can be generated by chemically cross-linking the CDK-binding motifs.

[0010] In preferred embodiments, at least one of the CDK-binding motifs is a CDK-binding motif of a CDK inhibitor protein, such as an INK4 protein, e.g., p15, p16, p18 and p19, or a CIP protein, e.g., p21^{CIP1}, p27^{KIP1}, and p57^{KIP2}. However, it will be understood that other CDK-binding motifs may be useful. Indeed, the CDK-binding motif of the INK4 proteins is characterized by tandemly arranged ankyrin-like sequences, which sequences exist in other proteins and, for those which are able to bind a CDK, can be used to generate the subject chimeric proteins. Likewise, the CDK-binding motif can be a p21/p27 inhibitory domain of a protein which has some homology with the CIP protein family. Exemplary chimeric proteins of the present invention are designated by SEQ ID No. 2, 5 and 7, and are encoded by the CDS's designated in SEQ ID No. 1, 4 and 6.

[0011] In preferred embodiments, the CDK-binding motifs of the chimeric protein have different binding specificities, relative to one and other, for cyclin dependent kinases. For instance, the chimeric protein can be generated with a CDK-binding motif from a protein which binds to and inhibits a CDK involved in progression of the cell cycle in G₀ and/or G₁ phase, and another CDK-binding motif from a protein which binds to and inhibits a CDK involved in progression of the cell cycle in S, G₂ and/or M phase. That is, the chimeric protein will bind to and inhibit a plurality (two or more) of cyclin dependent kinases which are active in different phases of the cell-cycle.

[0012] In most embodiments, the nucleic acid will further include a transcriptional regulatory sequence for controlling transcription of the nucleotide sequence encoding the chimeric polypeptide, e.g., the transcriptional regulatory sequence is operably linked to a chimeric gene encoding the chimeric polypeptide. For example, the present invention specifically contemplates recombinant transfection systems which include: (i) a gene construct including a nucleic acid encoding a chimeric polypeptide comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases, and operably linked to a transcriptional regulatory sequence for causing expression of the chimeric polypeptide in eukaryotic cells, and (ii) a gene delivery composition for delivering the gene construct to a cell and causing the cell to be transfected with the gene construct. For example, the gene construct can be derived from a viral vector, such as an adenoviral vector, an adeno-associated viral vector or a retroviral vector. In such embodiments, the gene delivery composition comprises a recombinant viral particle. In other embodiments, the gene construct can be delivered by such means as a liposome or a poly-cationic nucleic acid binding agent. For in vivo delivery to a mammal, such as a human, the gene delivery composition will further include a pharmaceutically acceptable carrier for administration to an animal, and, as necessary, will be a sterile preparation and substantially free of pyrogenic agents.

[0013] The present invention also pertains to preparations of such chimeric polypeptides. e.g., polypeptides which are generated from CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases. In preferred embodiments, the chimeric polypeptide is formulated in pharmaceutically acceptable carrier for delivery to a mammal. For example, the chimeric polypeptide can be formulated in liposomal preparations.

[0014] Still another aspect of the present invention related to transgenic animals which have cells harboring a nucleic acid one of the subject fusion proteins.

[0015] Yet another aspect of the present invention relates to recombinant transfection systems, comprising

[0016] (i) a first gene construct comprising a coding sequence encoding an inhibitory polypeptide comprising at least one CDK-binding motif for binding and inhibiting activation of a cyclin dependent kinase (cdk), which coding sequence is operably linked to a transcriptional regulatory sequence for causing expression of the first polypeptide in eukaryotic cells,

[0017] (ii) a second gene construct comprising a coding sequence encoding an endothelialization polypeptide which promotes endothelialization, and

[0018] (ii) a gene delivery composition for delivering the gene constructs to a cell and causing the cell to be transfected with the gene construct.

[0019] In preferred embodiments, the CDK-binding motif is a CDK-binding motif of a CDK inhibitor protein, such as an INK4 protein (e.g., p15, p16, p18 or p19), or a CIP/KIP protein (e.g., p21^{CIP1}, p27^{KIP1}, and p57^{KIP2}). In other preferred embodiments, the CDK-binding motif comprises tandemly arranged ankyrin-like sequences, or a p21/p27 inhibitory domain.

[0020] The inhibitory polypeptide can be a fusion protein comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases.

[0021] In preferred embodiments, the endothelialization polypeptide stimulates endothelial cell proliferation and/or stimulates migration of endothelial cells to a wound site. For instance, the endothelialization polypeptide is selected from the group consisting of angiogenic basic fibroblast growth factors (bFGF), acid fibroblast growth factor (aFGF), vascular endothelial growth factor (VEGF), platelet derived growth factor (PDGF), vascular permeability growth factor (VPF), and transforming growth factor beta (TGF-β).

[0022] In preferred embodiments, the the first and second gene constructs are provided as part of a single vector, and can be provided as part of polycistronic message. Alternatively, the the first and second gene constructs are provided on separate vectors.

[0023] In certain embodiments, the gene construct comprises a viral vector, e.g., an adenoviral vector, an adeno-associated viral vector, or a retroviral vector.

[0024] In other embodiments, the gene construct is provided in a delivery composition, e.g., selected from the group consisting of a liposome and a poly-cationic nucleic acid binding agent.

[0025] The subject transfection systems can be used for treating an animal having a vascular wound characterized a breach of endothelial integrity and by excessive smooth muscle proliferation, by a method which includes administering the recombinant transfection system to the area of the vascular wound. In preferred embodiments, the subject transfection systems are used in the treatment of restenosis, and may be administered by catheter.

[0026] Still another aspect of the present invention provides a gene construct encoding a fusion protein comprising a therapeutic polypeptide sequence from an intracellular protein which alter a biological process of a cell upon intracellular localization of the fusion protein, and a transcellular polypeptide sequence for promoting transcytosis of the fusion protein across a cell surface membrane and into a cell.

[0027] In preferred embodiments, the transcellular fusion protein alters one or more such biological processes as proliferation, differentiation, cell death, gene expression, protein stability, calcium mobilization, ion permeability, phosphorylation of intracellular proteins, metabolism of inositol phosphates (IP₃ and the like, diacyl glycerides, etc), and metabolism of nucleosides (such as cAMP).

[0028] The transcellular fusion protein can, by virtue of its binding to a protein or nucleic acid in the targeted cell, alter

(inhibit or potentiate) protein-protein interactions or protein-nucleic acid interactions between proteins and nucleic acids endogenous to the cell.

[0029] In certain embodiments, the therapeutic polypeptide sequence can be derived from a tumor suppressor, a transcription factor, a signal transduction protein, an antiviral protein or a metal chelating protein. For instance, the therapeutic polypeptide sequence can include a polypeptide sequence of a tumor suppressor such as p53, Rb or an Rb-like protein, or a CKI protein. In other embodiments, the therapeutic polypeptide sequence includes a polypeptide sequence of a signal transduction protein, such as from tubby, a DOT protein, a Bcl protein (bcl-2, bcl-x, etc), or an IκB protein.

[0030] In a preferred embodiment, the fusion protein includes at least one CDK-binding motif for binding and inhibiting activation of a cyclin dependent kinase (cdk). For instance, the CDK-binding motif is a CDK-binding motif of a CDK inhibitor protein, such as an INK4 protein (e.g., p15, p16, p18 or p19), or a CIP/KIP protein (e.g., p21^{CIP1}, p27^{KIP1}, and p57^{KIP2}). In other preferred embodiments, the CDK-binding motif comprises tandemly arranged ankyrin-like sequences, or a p21/p27 inhibitory domain. In preferred embodiments, the polypeptide comprises CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases.

[0031] The transcellular polypeptide sequence can be an internalizing peptide, such as may be derived from a polypeptide selected from the group consisting of antepennepedia protein, HIV transactivating (TAT) protein, mastoparan, melittin, bombolitin, delta hemolysin, pardaxin, Pseudomonas exotoxin A, clathrin, Diphtheria toxin and C9 complement protein.

[0032] In other embodiments, the transcellular polypeptide sequence can be an accessory peptide sequence which enhances interaction of the fusion protein with a cell surface membrane, such as a peptide sequence that includes an RGD sequence.

[0033] In preferred embodiments, the gene construct comprises a viral vector, e.g., an adenoviral vector, an adeno-associated viral vector, or a retroviral vector.

[0034] In other embodiments, the gene construct is provided in a delivery composition, e.g., selected from the group consisting of a liposome and a poly-cationic nucleic acid binding agent.

[0035] The invention also provides compositions of the fusion protein, e.g., such as may be formulated in pharmaceutical preparations.

[0036] The gene construct and the fusion protein may each be used as part of a method for treating an animal for unwanted cell proliferation, by administering the chimeric gene or the fusion protein.

[0037] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989);

DNA Cloning, Volumes I and II (D. N. Glover ed., 1985); *Oligonucleotide Synthesis* (M. J. Gait ed., 1984); Mullis et al. U.S. Pat. No: 4,683,195; *Nucleic Acid Hybridization* (B. D. Hames & S. J. Higgins eds. 1984); *Transcription And Translation* (B. D. Hames & S. J. Higgins eds. 1984); *Culture Of Animal Cells* (R. I. Freshney, Alan R. Liss, Inc., 1987); *Immobilized Cells And Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide To Molecular Cloning* (1984); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Gene Transfer Vectors For Mammalian Cells* (J. H. Miller and M. P. Calos eds., 1987, Cold Spring Harbor Laboratory); *Methods In Enzymology*, Vols. 154 and 155 (Wu et al. eds.), *Immunochemical Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London, 1987); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986); *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986).

DETAILED DESCRIPTION OF THE INVENTION

[0038] Progression of eukaryotic cells through the cell cycle is governed by the sequential formation, activation, and subsequent inactivation of a series of cyclin/cyclin dependent kinase complexes. The mechanisms underlying the expression of cyclins and the activation of the different cyclin-CDK complexes needed for progression through successive cell cycle transitions are now fairly well understood. In addition to positive regulation by the activation of cyclin-CDK complexes, negative regulation of the cell cycle occurs at checkpoints, many of which operate to control formation of cyclin/CDK complexes and/or activation of the complexes. Accordingly, these transitions are negatively regulated by signals that constrain the cell-cycle until specific conditions are fulfilled. Entry in to mitosis, for example, is inhibited by incompletely replicated DNA or DNA damage. These restriction on cell-cycle progression are essential for preserving the fidelity of the genetic information during cell division. The transition from G₁ to S phase, on the other hand, coordinates cell proliferation with environmental cues, after which the checks on the cell-cycle progression tend to be cell autonomous. Disruption of these signaling pathways can uncouple cellular responses from environmental controls and may lead to unrestrained cell proliferation or aberrant loss of differentiation.

[0039] (i) Overview

[0040] One aspect of the present invention pertains to novel inhibitors of cyclin-dependent kinases (CDKs), particularly CDK/cyclin complexes, which inhibitors can be used to control proliferation and/or differentiation of cells in which the inhibitors are introduced. More specifically, the inhibitors of the invention are chimeric proteins which include CDK-binding motifs from two or more different proteins. Such chimeric proteins are referred to herein as "poly-CBM proteins". For example, as set forth in greater detail below, the subject chimeric proteins can be generated from the in-frame fusion of coding sequences from two different CDK inhibitor proteins (generically referred to herein as "CKI" proteins), such as may be derived from fusion of coding sequences for an INK4 protein and coding sequences for a CIP protein. Moreover, as the appended examples describe, chimeric proteins of the present invention have been observed to be more potent inhibitors of

cyclin/CDK complexes than were either of the portions of the chimeric protein individually. For instance, p27-p16 chimeric proteins inhibited a cyclin D1/CDK4 complex with an IC_{50} more than two-fold less than p27 alone, and ten-fold less than p16 alone. Likewise, the p27-p16 chimeric protein inhibited cyclin E/CDK2, cyclin A/CDK2 and cyclin B/CDK2 complexes with IC_{50} 's approximately two-fold less than p27 alone (p16 itself not having any significant inhibitory activity against any of the three complexes).

[0041] Another aspect of the present invention pertains to chimeric polypeptides, particularly chimeric CKI polypeptides, which includes a heterologous peptide sequence which drives the translocation of an intracellular polypeptide across the cell membrane in order to facilitate localization of the polypeptide in the cellular cytoplasm and/or nucleoplasm to which the polypeptide is ectopically applied. Such chimeric polypeptides are referred to herein as "transcellular polypeptides". In general, such transcellular polypeptides includes a first peptide portion ("internalizing peptide") which by itself is capable of crossing the cellular membrane, e.g., by transcytosis, at a relatively high rate, or which facilitates such transcytosis. The internalizing peptide is conjugated to a polypeptide corresponding to a therapeutic polypeptide, e.g., having intracellular activity, such as the subject CKI polypeptides. The resulting chimeric polypeptide is transported into cells at a much higher rate than the therapeutic polypeptide alone would be, and thereby provide an effective means for introducing the therapeutic polypeptide into cells. The transcellular polypeptide can applied ectopically, or expressed and secreted by cells adjacent to the intended target cells. In the latter embodiment, the internalizing peptide facilitates the uptake of the therapeutic polypeptide by cells proximal to a cell which expresses the recombinant polypeptide. In this manner, the transcellular embodiment of the therapeutic polypeptide can be used to effectively treat cells which are, under the circumstances of treatment, e.g., inefficient or otherwise refractive to transfection with a gene construct encoding the therapeutic polypeptide. In a preferred embodiment, the transcellular polypeptide is delivered by way of a gene therapy expression vector—e.g., expression of the chimeric polypeptide by a subset of treated cells can result in localization of the polypeptide, with concomitant mitotic inhibition, in neighboring cells. For embodiments of a transcellular CKI polypeptide, the polypeptide can include all or a fragment of a single CKI protein, or include the poly-CBM protein described above.

[0042] In still another aspect, the present invention provides methods and compositions for enhancing endothelialization of injured sites while concurrently inhibiting smooth muscle cell proliferation around the site. In these embodiments, a CKI polypeptide is co-delivered with a polypeptide (an "endothelialization polypeptide") which stimulates (a) endothelial cell proliferation; (b) stimulates migration of endothelial cells to the wound site; (c) inhibits smooth muscle cell migration; and/or (d) inhibits proliferation of smooth muscle cells. In preferred embodiments, the CKI polypeptide and endothelialization polypeptide are ultimately delivered in the form of a single expression vector encoding both polypeptides, though multiple transfection of cells with separate expression vectors is also contemplated. The CKI polypeptide can include all or a fragment of a single CKI protein, or can be the poly-CBM protein described above. The co-delivery of the CKI polypeptide and endothelialization

polypeptide can have beneficial effects in, e.g., the repair of cardiovascular damage, repair of arteriosclerotic lesions and endothelialization of synthetic vascular grafts.

[0043] Other aspects of the present invention include: preparations of the subject polypeptides; expression constructs for recombinant production of the subject polypeptides, particularly for use as part of a gene therapy treatment; and methods for modulating cell proliferation and/or differentiation with the subject polypeptides.

[0044] (ii) Definitions

[0045] For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

[0046] The phrase "CDK-binding motif" refers to that portion of a protein which interacts either directly or indirectly with a cyclin dependent kinase (CDK). The binding motif may be a sequential portion of the protein, i.e., a contiguous sequence of amino acids, or it may be conformational, i.e. a combination of non-contiguous sequences of amino acids which, when the protein is in its native folding state, forms a structure which interacts with a CDK. The term "CDK-binding motif" explicitly includes any polypeptide which is identical, substantially homologous, or otherwise functionally or structurally equivalent to a portion of a CKI protein which binds directly or indirectly to a CDK or CDK complex. Other exemplary CDK-binding motifs can be provided from, for example, Rb and Rb-like proteins as well as cyclins.

[0047] An "inhibitor of CDK activation" refers to a molecule able to interact with a CDK and prevent activation of a kinase activity of the CDK either by, for example, inhibiting formation of CDK complexes including regulatory subunits, inhibiting interaction of the CDK subunit with activating kinases or phosphatases, inhibiting substrate binding, inhibiting ATP binding, and/or inhibiting conformational changes required for enzymatic activity. Accordingly, such inhibition may be by a direct, competitive mechanism, or by an indirect, non- or uncompetitive mechanism.

[0048] As used herein, the term "CKI protein" refers to a protein which can bind to and inhibit activation of a cyclin dependent kinase. Exemplary CKI proteins include members of the INK4 family, such as p16^{INK4A} or p15^{INK4B}, and members of the CIP family, such as p21^{CIP1}, p27^{KIP1}, and p57^{KIP2}.

[0049] The term "INK4 protein" refers to a family of structurally related CDK inhibitors characterized by a four-fold repeated ankyrin-like sequence (Elledge et al. (1994) *Curr. Opin. Cell Biol.* 6:874-878), and the ability to bind to CDKs, especially CDK4 and CDK6. Exemplary members of this protein family include p16 (INK4A/MTS1; Serrano et al. (1993) *Nature* 366:704-707); p15 (INK4B; Hannon et al. (1994) *Nature* 371:257-261); p18 (Guan et al. (1994) *Genes Dev.* 8:2939-2952) and p19 (Chan et al. (1995) *Mol. Cell Biol.* 15:2682-2688; and Hirai et al. (1995) *Mol. Cell Biol.* 15:2672-2681). Other proteins have been identified in the art as having tandemly arranged ankyrin-like sequences, such as the Pho81p protein (Ogawa et al. (1995) *Mol. Cell Biol.* 15:997-1004), and may provide CDK-binding motifs which are functionally equivalent to those of an INK4 protein.

[0050] The term "CIP protein" refers to members of another CKI protein family which includes p21^{CIP1} (WAF1/

SDI1/CAP20; Xiong et al. (1993) *Nature* 366:701-704); p27^{KIP1} (Polyak et al. (1994) *Cell* 78:67-74); and p57^{KIP2} (Lee et al. (1995) *Genes Dev.* 9:639-649; and Matsuoka et al. (1995) *Genes Dev.* 9:650-662). In addition to the functional characteristic of CDK inhibition, the CIP proteins each have a CDK inhibitory motif (a CDK-binding motif) of about 50 amino acids, referred to herein as a "p21/p27" inhibitory domain, which is conserved in members of the CIP family, as well as, for example, members of the Rb-like protein family.

[0051] A "chimeric protein" refers to a protein which includes polypeptide sequences from at least two different and distinct proteins. A chimeric protein can be a fusion protein, or the different polypeptide sequences can be covalently linked by a non-peptide bond, e.g., a cross-linking agent.

[0052] As used herein, the term "fusion protein" is art recognized and refer to a chimeric protein which is at least initially expressed as single chain protein comprised of amino acid sequences derived from two or more different proteins, e.g., the fusion protein is a gene product of a fusion gene.

[0053] The art term "fusion gene" refers to a nucleic acid in which two or more genes are fused resulting in a single open reading frame for coding two or more proteins that as a result of this fusion are joined by one or more peptide bonds.

[0054] As used herein, the term "nucleic acid" refers to polynucleotides such as deoxyribonucleic acid (DNA), and, where appropriate, ribonucleic acid (RNA). The term should also be understood to include, as equivalents, analogs of either RNA or DNA made from nucleotide analogs, and, as applicable to the embodiment being described, single-stranded (such as sense or antisense) and double-stranded polynucleotides.

[0055] As used herein, the term "gene" or "recombinant gene" refers to a nucleic acid comprising an open reading frame encoding a fusion polypeptide of the present invention, including both exonic and (optionally) intronic sequences. An exemplary recombinant gene encoding a subject fusion protein is represented by SEQ. ID No: 1.

[0056] As used herein, the term "transfection" means the introduction of a heterologous nucleic acid, e.g., an expression vector, into a recipient cell by nucleic acid-mediated gene transfer. "Transformation", as used herein with respect to transfected nucleic acid, refers to a process in which a cell's genotype is changed as a result of the cellular uptake of exogenous DNA or RNA, and, for example, the transformed cell expresses a recombinant form of a CDK-inhibitory fusion polypeptide of the present invention.

[0057] "Expression vector" refers to a replicable DNA construct used to express DNA which encodes the desired protein and which includes a transcriptional unit comprising an assembly of (1) genetic element(s) having a regulatory role in gene expression, for example, promoters, operators, or enhancers, operatively linked to (2) a DNA sequence encoding a desired protein (in this case, a fusion protein of the present invention) which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences. The choice of promoter and other regulatory elements generally varies

according to the intended host cell. In general, expression vectors of utility in recombinant DNA techniques are often in the form of "plasmids" which refer to circular double stranded DNA loops which, in their vector form are not bound to the chromosome. In the present specification, "plasmid" and "vector" are used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which become known in the art subsequently hereto.

[0058] In the expression vectors, regulatory elements controlling transcription or translation can be generally derived from mammalian, microbial, viral or insect genes. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants may additionally be incorporated. Vectors derived from viruses, such as retroviruses, adenoviruses, and the like, may be employed.

[0059] "Transcriptional regulatory sequence" is a generic term used throughout the specification to refer to DNA sequences, such as initiation signals, enhancers, and promoters and the like which induce or control transcription of protein coding sequences with which they are operably linked. In preferred embodiments, transcription of the fusion gene is under the control of a promoter sequence (or other transcriptional regulatory sequence) which controls the expression of the recombinant gene in a cell-type in which expression is intended. It will also be understood that the recombinant gene can be under the control of transcriptional regulatory sequences which are the same or which are different from those sequences which control transcription of one of the naturally-occurring forms of a CDK inhibitor protein.

[0060] As used herein, the term "tissue-specific promoter" means a DNA sequence that serves as a promoter, i.e., regulates expression of a selected DNA sequence operably linked to the promoter, and which effects expression of the selected DNA sequence in specific cells of a tissue, such as cells of a urogenital origin, e.g. renal cells, or cells of a neural origin, e.g. neuronal cells. The term also covers so-called "leaky" promoters, which regulate expression of a selected DNA primarily in one tissue, but cause expression in other tissues as well.

[0061] "Operably linked" when describing the relationship between two DNA regions simply means that they are functionally related to each other. For example, a promoter or other transcriptional regulatory sequence is operably linked to a coding sequence if it controls the transcription of the coding sequence.

[0062] "Recombinant host cells" refers to cells which have been transformed or transfected with vectors constructed using recombinant DNA techniques. As relevant to the present invention, recombinant host cells are those which produce CDK inhibitor fusion proteins by virtue of having been transformed with expression vectors encoding these proteins.

[0063] As used herein, a "transgenic animal" is any animal, preferably a non-human mammal, a bird or an amphibian, in which one or more of the cells of the animal contain heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in

the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by micro-injection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. This molecule may be integrated within a chromosome, or it may be extrachromosomally replicating DNA. In the typical transgenic animals described herein, the transgene causes cells to express a recombinant form of a CDK inhibitory fusion protein. The term "chimeric animal" is used herein to refer to animals in which the recombinant gene is found, or in which the recombinant is expressed in some but not all cells of the animal. The term "tissue-specific chimeric animal" indicates that the recombinant gene is present and/or expressed in some tissues but not others.

[0064] "Homology" refers to sequence similarity between two peptides or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same base or amino acid, then the molecules are homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences.

[0065] (iii) Poly-CBM Polypeptides

[0066] One aspect of the invention pertains to chimeric proteins which include CDK-binding motifs from two or more different proteins. In one embodiment, the invention provides a nucleic acid having a nucleotide sequence encoding a chimeric CDK inhibitor protein, and/or equivalents of such nucleic acids. In general, the nucleic acid is derived by the in-frame fusion of coding sequences from two or more proteins which have CDK-inhibitory motifs, such motifs being preserved in the resultant chimeric protein. Accordingly, such chimeric proteins can be derived to include, for example, CKI protein sequences, such as from INK4 or CIP proteins. For instance, as described in the appended examples, a coding sequence providing the CDK-binding motif of an INK4 protein can be fused in frame to a coding sequence providing a CDK-binding motif of a CIP protein.

[0067] Exemplary nucleic acid of the present invention encode fusion proteins which include at least a CDK-binding portion of an INK4 protein, such as p15, p16, p18 or p19. In preferred embodiments, the chimeric protein includes at least the two ankyrin-like sequence of the C-terminal portion of the INK4 protein, e.g. corresponding to the 3rd repeat (residues 69-101) and 4th repeat (residues 102-133) of p16^{INK4A} (see Serrano et al. (1993) Nature 366:704-707).

[0068] Similarly, preferred chimeric proteins of the present invention include at least the p21/27-related inhibitory domain of a CIP protein, e.g. from p21, p27 or p57. For example, the chimeric protein can include the CDK-inhibitory motif corresponding to residues 28-79 of p27, residues 17-68 of p21, and/or residues 31-82 of p57, though larger fragments may be used such as described in the appended examples.

[0069] Moreover, CDK-binding motifs homologous to those occurring in either the INK4 or CIP protein families

have been observed in other proteins. For example, the p21/p27-related inhibitory domain typical of the CIP protein family has been identified in such other proteins as the Rb-related protein p107 (Zhu et al. (1995) Genes Dev 9:1740-1752). Likewise, ankyrin-like repeats homologous with the INK4 proteins have been identified in such other proteins as the Pho81p protein (Ogawa et al. (1995) Mol Cell Biol 15:997-1004). Consequently, it will be apparent to one of ordinary skill in the art, based on the disclosure herein, that functional equivalents of the INK4 and CIP proteins, e.g. which are capable of binding to a CDK and inhibiting kinase activation, exist and can be provided in the subject chimeric proteins.

[0070] Furthermore, it will be understood that the subject chimeric proteins can include CDK-binding motifs from proteins unrelated to either the INK4 family or CIP family. Moreover, such CDK-binding motifs, while inhibitory in and of themselves, can be derived from proteins which are otherwise activating in their full length form. To illustrate, the subject chimeric protein can be generated with a fragment of a cyclin which retains its CDK binding ability but not the CDK activating ability characteristic of the full length protein.

[0071] In some instances it may be necessary to introduce an unstructured polypeptide linker region between portions of the chimeric protein derived from different proteins. This linker can facilitate enhanced flexibility of the chimeric protein allowing the CDK-binding motifs from each portion to freely and (optionally) simultaneously interact with a CDK by reducing steric hindrance between the two fragments, as well as allowing appropriate folding of each portion to occur. The linker can be of natural origin, such as a sequence determined to exist in random coil between two domains of a protein. Alternatively, the linker can be of synthetic origin. For instance, the sequence (Gly,Ser)₃ can be used as a synthetic unstructured linker. Linkers of this type are described in Huston et al. (1988) PNAS 85:4879; and U.S. Pat. Nos. 5,091,513 and 5,258,498. Naturally occurring unstructured linkers of human origin are preferred as they reduce the risk of immunogenicity.

[0072] Techniques for making fusion genes are well known. Essentially, the joining of various DNA fragments coding for different polypeptide sequences is performed in accordance with conventional techniques, employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed to generate a fusion gene sequence (see, for example, Current Protocols in Molecular Biology, eds. Ausubel et al. John Wiley & Sons: 1992).

[0073] The term nucleic acid as used herein is intended to include nucleotide sequences encoding functionally equivalent chimeric proteins which, for example, retain the ability to bind to a cyclin-dependent kinase. Equivalent nucleotide sequences will include sequences that differ by one or more

nucleotide substitutions, additions or deletions, such as allelic variants; and will, therefore, include sequences that differ from the nucleotide sequence of, for example, an INK4 or CIP gene known in the art due to the degeneracy of the genetic code. Equivalents will also include nucleotide sequences that hybridize under stringent conditions (i.e., equivalent to about 20-27° C. below the melting temperature (T_m) of the DNA duplex formed in about 1M salt) to the nucleotide sequence encoding a naturally-occurring CDK-binding motif. Furthermore, equivalent nucleic acids will include those with nucleotide sequences which differ from the natural sequence which encodes a CDK-binding motif because of degeneracy in the genetic code. For example, a number of amino acids are designated by more than one triplet. Codons that specify the same amino acid can, accordingly, be used to replace codons in the naturally-occurring sequence.

[0074] This invention also provides expression vectors comprising a nucleotide sequence encoding a subject CDK inhibitor chimeric protein and operably linked to at least one regulatory sequence. Regulatory sequences are art-recognized and are selected to direct expression of the fusion protein. Accordingly, the term regulatory sequence includes promoters, enhancers and other expression control elements. Exemplary regulatory sequences are described in Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990). For instance, any of a wide variety of expression control sequences—sequences that control the expression of a DNA sequence when operatively linked to it may be used in these vectors to express DNA sequences encoding the fusion proteins of this invention. Such useful expression control sequences, include, for example, the early and late promoters of SV40, adenovirus or cytomegalovirus immediate early promoter, the lac system, the trp system, the TAC or TRC system, T7 promoter whose expression is directed by T7 RNA polymerase, the major operator and promoter regions of phage lambda, the control regions for fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast α -mating factors, the polyhedron promoter of the baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. Of course, the transcriptional regulatory sequences can include those sequences which naturally control expression of one of the genes used to derive the fusion protein, such as 5' flanking sequences of an INK4 or CIP gene.

[0075] It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed. Moreover, the vector's copy number, the ability to control that copy number and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered.

[0076] Expression vehicles for production of recombinant forms of the subject chimeric proteins include plasmids and other vectors. For instance, suitable vectors for expression of a fusion protein of the present invention include plasmids of the types: pBR322-derived plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli*.

[0077] A number of vectors exist for the expression of recombinant proteins in yeast. For instance, YEP24, YIP5, YEP51, YEP52, pYES2, and YRP17 are cloning and expression vehicles useful in the introduction of genetic constructs into *S. cerevisiae* (see, for example, Broach et al. (1983) in Experimental Manipulation of Gene Expression, ed. M. Inouye Academic Press, p. 83). These vectors can replicate in *E. coli* due the presence of the pBR322 ori, and in *S. cerevisiae* due to the replication determinant of the yeast 2 micron plasmid. In addition, drug resistance markers such as ampicillin can be used.

[0078] The preferred mammalian expression vectors (other than for gene therapy) contain both prokaryotic sequences to facilitate the propagation of the vector in bacteria, and one or more eukaryotic transcription units that are expressed in eukaryotic cells. The pcDNA1/amp, pcDNA1/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. Alternatively, derivatives of viruses such as the bovine papilloma virus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins in eukaryotic cells.

[0079] In some instances, it may be desirable to express the subject fusion protein by the use of a baculovirus expression system. Examples of such baculovirus expression systems include pVL-derived vectors (such as pVL1392, pVL1393 and pVL941), pAcUW-derived vectors (such as pAcUW1), and pBlueBac-derived vectors (such as the β -gal containing pBlueBac III).

[0080] The various methods employed in the preparation of the plasmids and transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see Molecular Cloning: A Laboratory Manual, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989) Chapters 16 and 17.

[0081] Another aspect of the present invention concerns preparations of the subject chimeric proteins. In particular, purified and semi-purified preparations of the CDK inhibitors can be formulated according to specifications attendant the desired use of the chimeric protein.

[0082] With respect to purifying the subject chimeric proteins, Applicant notes that it is widely appreciated that addition of certain heterologous sequences to a protein can facilitate the expression and purification of the proteins. For example, a fusion protein of the present invention can be generated to also include a glutathione-S-transferase (GST) polypeptide sequence. The GST portion of the recombinant proteins can enable easy purification of the protein, such as by the use of glutathione-derivatized matrices (see, for example, Current Protocols in Molecular Biology, eds. Ausabel et al. (N.Y.: John Wiley & Sons, 1991)). In another embodiment, the subject fusion protein can also include a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence located at the N-terminus of the subject fusion protein. Such sequences facilitates purifica-

tion of the poly(His)-expressed fusion protein by affinity chromatography using a Ni^{2+} -metal resin. The purification leader sequence can then be subsequently removed by treatment with enterokinase (e.g., see Hochuli et al. (1987) *J. Chromatography* 411:177; and Janknecht et al. *PNAS* 88:8972).

[0083] The present invention further pertains to methods of producing the subject chimeric proteins. For example, a host cell transfected with a nucleic acid vector directing expression of a nucleotide sequence encoding one of the chimeric proteins of the present invention can be cultured under appropriate conditions to allow expression of the polypeptide to occur. The peptide may be secreted and isolated from a mixture of host cells and medium by inclusion of a signal secretion sequence. Alternatively, the peptide may be retained cytoplasmically and the cells harvested, lysed and the protein isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. The recombinant chimeric protein can be isolated from cell culture medium, host cells, or both using techniques known in the art for purifying proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immuno-affinity purification with antibodies specific for portions of the chimeric protein.

[0084] This invention also pertains to a host cell transfected to recombinantly express one of the subject chimeric proteins. The host cell may be any prokaryotic or eukaryotic cell. Thus, a nucleic acid derived from the fusion of coding sequences for two or more CDK-binding motifs from different proteins can be used to produce a recombinant form of the chimeric protein via microbial or eukaryotic cellular processes. Ligating the polynucleotide sequence into a gene construct, such as an expression vector, and transforming or transfecting into hosts, either eukaryotic (yeast, avian, insect or mammalian) or prokaryotic (bacterial cells), are standard procedures used in producing other well-known proteins, e.g., p16, p21, p27, p57, p107, cyclins and the like. Similar procedures, or modifications thereof, can be employed to prepare recombinant chimeric proteins by microbial means or tissue-culture technology in accord with the subject invention.

[0085] The chimeric molecules of the present invention can also be generated using well-known cross-linking reagents and protocols. For example, there are a large number of chemical cross-linking agents that are known to those skilled in the art and useful for cross-linking two heterologous polypeptide chains. For the present invention, the preferred cross-linking agents are heterobifunctional cross-linkers, which can be used to link molecules in a stepwise manner. Heterobifunctional cross-linkers provide the ability to design more specific coupling methods for conjugating proteins, thereby reducing the occurrences of unwanted side reactions such as homo-protein polymers. A wide variety of heterobifunctional cross-linkers are known in the art. These include: succinimidyl 4-(N-maleimidomethyl)cyclohexane 1-carboxylate (SMCC), m-Maleimido-benzoyl-N-hydroxysuccinimide ester (MBS); N-succinimidyl(4-iodoacetyl)aminobenzoate (SIAB), succinimidyl 4-(p-maleimidophenyl)butyrate (SMPB), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC); 4-succinimidylloxycarbonyl-a-methyl-a-(2-pyridyldithio)-toluene (SMPT), N-succinimidyl 3-(2-pyridyldithio) propi-

onate (SPDP), succinimidyl 6-[3-(2-pyridyldithio)propionate]hexanoate (LC-SPDP). Those cross-linking agents having N-hydroxysuccinimide moieties can be obtained as the N-hydroxysulfosuccinimide analogs, which generally have greater water solubility. In addition, those cross-linking agents having disulfide bridges within the linking chain can be synthesized instead as the alkyl derivatives so as to reduce the amount of linker cleavage in vivo.

[0086] In addition to the heterobifunctional cross-linkers, there exists a number of other cross-linking agents including homobifunctional and photoreactive cross-linkers. Disuccinimidyl suberate (DSS), bismaleimido-hexane (BMH) and dimethylpimelimidate.2 HCl (DMP) are examples of useful homobifunctional cross-linking agents, and bis-[β -(4-azidosalicylamido)ethyl]disulfide (BASED) and N-succinimidyl-6-(4'-azido-2'-nitrophenylamino)hexanoate (SANPAH) are examples of useful photoreactive cross-linkers for use in this invention. For a recent review of protein coupling techniques, see Means et al. (1990) *Bioconjugate Chemistry* 1:2-12, incorporated by reference herein.

[0087] One particularly useful class of heterobifunctional cross-linkers, included above, contain the primary amine reactive group, N-hydroxysuccinimide (NHS), or its water soluble analog N-hydroxysulfosuccinimide (sulfo-NHS). Primary amines (lysine epsilon groups) at alkaline pH's are unprotonated and react by nucleophilic attack on NHS or sulfo-NHS esters. This reaction results in the formation of an amide bond, and release of NHS or sulfo-NHS as a by-product.

[0088] Another reactive group useful as part of a heterobifunctional cross-linker is a thiol reactive group. Common thiol reactive groups include maleimides, halogens, and pyridyl disulfides. Maleimides react specifically with free sulfhydryls (cysteine residues) in minutes, under slightly acidic to neutral (pH 6.5-7.5) conditions. Halogens (iodoacetyl functions) react with —SH groups at physiological pH's. Both of these reactive groups result in the formation of stable thioether bonds.

[0089] The third component of the heterobifunctional cross-linker is the spacer arm or bridge. The bridge is the structure that connects the two reactive ends. The most apparent attribute of the bridge is its effect on steric hindrance. In some instances, a longer bridge can more easily span the distance necessary to link two complex biomolecules. For instance, SMPB has a span of 14.5 angstroms.

[0090] Preparing protein-conjugates using heterobifunctional reagents is a two-step process involving the amine reaction and the sulfhydryl reaction. For the first step, the amine reaction, the protein chosen should contain a primary amine. This can be lysine epsilon amines or a primary alpha amine found at the N-terminus of most proteins. The protein should not contain free sulfhydryl groups. In cases where both proteins to be conjugated contain free sulfhydryl groups, one protein can be modified so that all sulfhydryls are blocked using for instance, N-ethylmaleimide (see Partis et al. (1983) *J. Pro. Chem.* 2:263, incorporated by reference herein). Ellman's Reagent can be used to calculate the quantity of sulfhydryls in a particular protein (see for example Ellman et al. (1958) *Arch. Biochem. Biophys.* 74:443 and Riddles et al. (1979) *Anal. Biochem.* 94:75, incorporated by reference herein).

[0091] The reaction buffer should be free of extraneous amines and sulfhydryls. The pH of the reaction buffer should

be 7.0-7.5. This pH range prevents maleimide groups from reacting with amines, preserving the maleimide group for the second reaction with sulfhydryls.

[0092] The NHS-ester containing cross-linkers have limited water solubility. They should be dissolved in a minimal amount of organic solvent (DMF or DMSO) before introducing the cross-linker into the reaction mixture. The cross-linker/solvent forms an emulsion which will allow the reaction to occur.

[0093] The sulfo-NHS ester analogs are more water soluble, and can be added directly to the reaction buffer. Buffers of high ionic strength should be avoided, as they have a tendency to "salt out" the sulfo-NHS esters. To avoid loss of reactivity due to hydrolysis, the cross-linker is added to the reaction mixture immediately after dissolving the protein solution.

[0094] The reactions can be more efficient in concentrated protein solutions. The more alkaline the pH of the reaction mixture, the faster the rate of reaction. The rate of hydrolysis of the NHS and sulfo-NHS esters will also increase with increasing pH. Higher temperatures will increase the reaction rates for both hydrolysis and acylation.

[0095] Once the reaction is completed, the first protein is now activated, with a sulfhydryl reactive moiety. The activated protein may be isolated from the reaction mixture by simple gel filtration or dialysis. To carry out the second step of the cross-linking, the sulfhydryl reaction, the protein chosen for reaction with maleimides, activated halogens, or pyridyl disulfides must contain a free sulfhydryl, usually from a cysteine residue. Free sulfhydryls can be generated by reduction of protein disulfides. Alternatively, a primary amine may be modified with Traut's Reagent to add a sulfhydryl (Blattler et al. (1985) *Biochem* 24:1517, incorporated by reference herein). Again, Ellman's Reagent can be used to calculate the number of sulfhydryls available in protein.

[0096] In all cases, the buffer should be degassed to prevent oxidation of sulfhydryl groups. EDTA may be added to chelate any oxidizing metals that may be present in the buffer. Buffers should be free of any sulfhydryl containing compounds.

[0097] Maleimides react specifically with —SH groups at slightly acidic to neutral pH ranges (6.5-7.5). A neutral pH is sufficient for reactions involving halogens and pyridyl disulfides. Under these conditions, maleimides generally react with —SH groups within a matter of minutes. Longer reaction times are required for halogens and pyridyl disulfides.

[0098] The first sulfhydryl reactive-protein prepared in the amine reaction step is mixed with the sulfhydryl-containing protein under the appropriate buffer conditions. The protein-protein conjugates can be isolated from the reaction mixture by methods such as gel filtration or by dialysis.

[0099] (iv) Transcellular Therapeutic Polypeptides

[0100] Another aspect of the present invention pertains to chimeric polypeptides which includes a heterologous peptide sequence ("internalizing peptide") which drives the translocation of an extracellular form of a therapeutic polypeptide sequence across a cell membrane in order to facilitate intracellular localization of the therapeutic

polypeptide. In this regard, the therapeutic polypeptide sequence is one which is active intracellularly, such as a tumor suppressor polypeptide, transcription factor or the like. The internalizing peptide, by itself, is capable of crossing a cellular membrane by, e.g., transcytosis, at a relatively high rate. The internalizing peptide is conjugated, e.g., as a fusion protein, to a therapeutic polypeptide. The resulting chimeric polypeptide is transported into cells at a higher rate relative to the polypeptide alone to thereby provide an means for enhancing the introduction of inhibitory polypeptides into surrounding cells, e.g., to enhance gene therapy and/or topical applications of the therapeutic polypeptide. For convenience, the transcellular therapeutic polypeptides are described below as fusion proteins including CKI polypeptide sequences, though as also described below (section v), many other protein domains can be used in place of the CKI polypeptide.

[0101] In one embodiment, the internalizing peptide is derived from the drosopholia antepennepedia protein, or homologs thereof. The 60 amino acid long long homeodomain of the homeo-protein antepennepedia has been demonstrated to translocate through biological membranes and can facilitate the translocation of heterologous polypeptides to which it is couples. See for example Derossi et al. (1994) *J Biol Chem* 269:10444-10450; and Perez et al. (1992) *J Cell Sci* 102:717-722. Recently, it has been demonstrated that fragments as small as 16 amino acids long of this protein are sufficient to drive internalization. See Derossi et al. (1996) *J Biol Chem* 271:18188-18193. The present invention contemplates a chimeric protein comprising at least one CDK binding motif and at least a portion of the antepennepedia protein (or homolog thereof) sufficient to increase the transmembrane transport of the chimeric protein, relative to the CDK binding motif alone, by a statistically significant amount.

[0102] Another example of an internalizing peptide is the HIV transactivator (TAT) protein. This protein appears to be divided into four domains (Kuppuswamy et al. (1989) *Nucl. Acids Res.* 17:3551-3561). Purified TAT protein is taken up by cells in tissue culture (Frankel and Pabo, (1989) *Cell* 55:1189-1193), and peptides, such as the fragment corresponding to residues 37-62 of TAT, are rapidly taken up by cell in vitro (Green and Loewenstein, (1989) *Cell* 55:1179-1188). The highly basic region mediates internalization and targeting of the internalizing moiety to the nucleus (Ruben et al., (1989) *J. Virol.* 63:1-8). Peptides or analogs that include a sequence present in the highly basic region, such as CFITKALGISYGRKKRRQRRPPQGS, are conjugated to CKI polypeptides (or CDK binding motifs thereof) to aid in internalization and targeting those proteins to the intracellular milieu.

[0103] Another exemplary transcellular CKI polypeptide can be generated to include a sufficient portion of mastoparan (T. Higashijima et al., (1990) *J. Biol. Chem.* 265:14176) to increase the transmembrane transport of the chimeric protein.

[0104] While not wishing to be bound by any particular theory, it is noted that hydrophilic polypeptides may be also be physiologically transported across the membrane barriers by coupling or conjugating the polypeptide to a transportable peptide which is capable of crossing the membrane by receptor-mediated transcytosis. Suitable internalizing pep-

tides of this type can be generated using all or a portion of, e.g., a histone, insulin, transferrin, basic albumin, prolactin and insulin-like growth factor I (IGF-I), insulin-like growth factor II (IGF-II) or other growth factors. For instance, it has been found that an insulin fragment, showing affinity for the insulin receptor on capillary cells, and being less effective than insulin in blood sugar reduction, is capable of transmembrane transport by receptor-mediated transcytosis and can therefor serve as an internalizing peptide for the subject transcellular CKI polypeptides. Preferred growth factor-derived internalizing peptides include EGF (epidermal growth factor)-derived peptides, such as CMHIESLDSYTC and CMYIEALDKYAC; TGF-beta (transforming growth factor beta)-derived peptides; peptides derived from PDGF (platelet-derived growth factor) or PDGF-2; peptides derived from IGF-I (insulin-like growth factor) or IGF-II; and FGF (fibroblast growth factor)-derived peptides.

[0105] Another class of translocating/internalizing peptides exhibits pH-dependent membrane binding. For an internalizing peptide that assumes a helical conformation at an acidic pH, the internalizing peptide acquires the property of amphiphilicity, e.g., it has both hydrophobic and hydrophilic interfaces. More specifically, within a pH range of approximately 5.0-5.5, an internalizing peptide forms an alpha-helical, amphiphilic structure that facilitates insertion of the moiety into a target membrane. An alpha-helix-inducing acidic pH environment may be found, for example, in the low pH environment present within cellular endosomes. Such internalizing peptides can be used to facilitate transport of CKI polypeptides, taken up by an endocytic mechanism, from endosomal compartments to the cytoplasm.

[0106] A preferred pH-dependent membrane-binding internalizing peptide includes a high percentage of helix-forming residues, such as glutamate, methionine, alanine and leucine. In addition, a preferred internalizing peptide sequence includes ionizable residues having pKa's within the range of pH 5-7, so that a sufficient uncharged membrane-binding domain will be present within the peptide at pH 5 to allow insertion into the target cell membrane.

[0107] A particularly preferred pH-dependent membrane-binding internalizing peptide in this regard is aa1-aa2-aa3-EAALA(EALA)4-EALEALAA-amide, which represents a modification of the peptide sequence of Subbarao et al. (*Biochemistry* 26:2964, 1987). Within this peptide sequence, the first amino acid residue (aa1) is preferably a unique residue, such as cysteine or lysine, that facilitates chemical conjugation of the internalizing peptide to a targeting protein conjugate. Amino acid residues 2-3 may be selected to modulate the affinity of the internalizing peptide for different membranes. For instance, if both residues 2 and 3 are lys or arg, the internalizing peptide will have the capacity to bind to membranes or patches of lipids having a negative surface charge. If residues 2-3 are neutral amino acids, the internalizing peptide will insert into neutral membranes.

[0108] Yet other preferred internalizing peptides include peptides of apo-lipoprotein A-1 and B; peptide toxins, such as melittin, bombolittin, delta hemolysin and the pardaxins; antibiotic peptides, such as alamethicin; peptide hormones, such as calcitonin, corticotrophin releasing factor, beta endorphin, glucagon, parathyroid hormone, pancreatic polypeptide; and peptides corresponding to signal sequences

of numerous secreted proteins. In addition, exemplary internalizing peptides may be modified through attachment of substituents that enhance the alpha-helical character of the internalizing peptide at acidic pH.

[0109] Yet another class of internalizing peptides suitable for use within the present invention include hydrophobic domains that are "hidden" at physiological pH, but are exposed in the low pH environment of the target cell endosome. Upon pH-induced unfolding and exposure of the hydrophobic domain, the moiety binds to lipid bilayers and effects translocation of the covalently linked CKI polypeptide into the cell cytoplasm. Such internalizing peptides may be modeled after sequences identified in, e.g., *Pseudomonas* exotoxin A, clathrin, or Diphtheria toxin.

[0110] Pore-forming proteins or peptides may also serve as internalizing peptides herein. Pore forming proteins or peptides may be obtained or derived from, for example, C9 complement protein, cytolytic T-cell molecules or NK-cell molecules. These moieties are capable of forming ring-like structures in membranes, thereby allowing transport of attached CKI polypeptide through the membrane and into the cell interior.

[0111] Mere membrane intercalation of an internalizing peptide may be sufficient for translocation of the CKI polypeptide across cell membranes. However, translocation may be improved by attaching to the internalizing peptide a substrate for intracellular enzymes (i.e., an "accessory peptide"). It is preferred that an accessory peptide be attached to a portion(s) of the internalizing peptide that protrudes through the cell membrane to the cytoplasmic face. The accessory peptide may be advantageously attached to one terminus of a translocating/internalizing moiety or anchoring peptide. An accessory moiety of the present invention may contain one or more amino acid residues. In one embodiment, an accessory moiety may provide a substrate for cellular phosphorylation (for instance, the accessory peptide may contain a tyrosine residue).

[0112] An exemplary accessory moiety in this regard would be a peptide substrate for N-myristoyl transferase, such as GNAAAARR (Eubanks et al., in *Peptides. Chemistry and Biology*, Garland Marshall (ed.), ESCOM, Leiden, 1988, pp. 566-69). In this construct, an internalizing, peptide would be attached to the C-terminus of the accessory peptide, since the N-terminal glycine is critical for the accessory moiety's activity. This hybrid peptide, upon attachment to a CKI polypeptide at its C-terminus, is N-myristylated and further anchored to the target cell membrane, e.g., it serves to increase the local concentration of the CKI polypeptide at the cell membrane.

[0113] To further illustrate use of an accessory peptide, a phosphorylatable accessory peptide is first covalently attached to the C-terminus of an internalizing peptide and then incorporated into a fusion protein with a CKI polypeptide. The peptide component of the fusion protein intercalates into the target cell plasma membrane and, as a result, the accessory peptide is translocated across the membrane and protrudes into the cytoplasm of the target cell. On the cytoplasmic side of the plasma membrane, the accessory peptide is phosphorylated by cellular kinases at neutral pH. Once phosphorylated, the accessory peptide acts to irreversibly anchor the fusion protein into the membrane. Localization to the cell surface membrane can enhance the translocation of the CKI polypeptide into the cell cytoplasm.

[0114] Suitable accessory peptides include peptides that are kinase substrates, peptides that possess a single positive charge, and peptides that contain sequences which are glycosylated by membrane-bound glycotransferases. Accessory peptides that are glycosylated by membrane-bound glycotransferases may include the sequence x-NLT-x, where "x" may be another peptide, an amino acid, coupling agent or hydrophobic molecule, for example. When this hydrophobic tripeptide is incubated with microsomal vesicles, it crosses vesicular membranes, is glycosylated on the luminal side, and is entrapped within the vesicles due to its hydrophilicity (C. Hirschberg et al., (1987) *Ann. Rev. Biochem.* 56:63-87). Accessory peptides that contain the sequence x-NLT-x thus will enhance target cell retention of corresponding CKI polypeptide.

[0115] In another embodiment of this aspect of the invention, an accessory peptide can be used to enhance interaction of the CKI polypeptide with the target cell. Exemplary accessory peptides in this regard include peptides derived from cell adhesion proteins containing the sequence "RGD", or peptides derived from laminin containing the sequence CDPGYIGSRC. Extracellular matrix glycoproteins, such as fibronectin and laminin, bind to cell surfaces through receptor-mediated processes. A tripeptide sequence, RGD, has been identified as necessary for binding to cell surface receptors. This sequence is present in fibronectin, vitronectin, C3bi of complement, von-Willebrand factor, EGF receptor, transforming growth factor beta, collagen type I, lambda receptor of *E. coli*, fibrinogen and Sindbis coat protein (E. Ruoslahti, *Ann. Rev. Biochem.* 57:375-413, 1988). Cell surface receptors that recognize RGD sequences have been grouped into a superfamily of related proteins designated "integrins". Binding of "RGD peptides" to cell surface integrins will promote cell-surface retention, and ultimately translocation, of the CKI integrin protein.

[0116] As described for the poly-CBM proteins above, the internalizing and accessory peptides can each, independently, be added to a CKI polypeptide by either chemical cross-linking or in the form of a fusion protein. In the

instance of fusion proteins, unstructured polypeptide linkers can be included between each of the peptide moieties.

[0117] The CKI polypeptide can consist of as little as a CDK-binding moiety of a CKI protein, or can include a full length CKI protein and/or poly-CBM protein.

[0118] In general, the internalization peptide will be sufficient to also direct export of the CKI polypeptide. However, where an accessory peptide is provided, such as an RGD sequence, it may be necessary to include a secretion signal sequence to direct export of the fusion protein from its host cell. In preferred embodiments, the secretion signal sequence is located at the extreme N-terminus, and is (optionally) flanked by a proteolytic site between the secretion signal and the rest of the fusion protein.

[0119] In an exemplary embodiment, the CKI polypeptides is engineered to include an integrin-binding RGD peptide/SV40 nuclear localization signal (see, for example Hart S L et al., 1994; *J. Biol. Chem.*, 269:12468-12474), such as encoded by the nucleotide sequence provided in the Nde1-EcoR1 fragment: catatgggtggctgcccgtggcgtatgttcggttgcggtgctcctccaaaaagaagagaaagtagctggattc, which encodes the RGD/SV40 nucleotide sequence: MGGCRGD-MFGCGAPPKKRKRVAGF. In another embodiment, the protein can be engineered with the HIV-1 tat(1-72) polypeptide, e.g., as provided by the Nde1-EcoR1 fragment:catatggccagtagatcctagactagagcccggaagcatccaggaagtcagcctaaactgcttgtaaccaattgctattgtaaaaagtgttgcctttcattgc-caagttgtttcatacaaaaagcccttg-gcatctcctatggcaggaagaagcg-gagacagcgacgaagacctcctcaaggcagtcagactcatcaagtttctctaagtaagcaagattc, which encodes the HIV-1 tat(1-72) peptide sequence: MEPVDPRLEPWKHPGSQPKTACTNCYCK-KCCFHCQVCFITKALGISYGRKKRRQR-RRPPQGSQTHQVSLSKQ. In still another embodiment, the fusion protein includes the HSV-1 VP22 polypeptide (Elliott G., O'Hare P (1997) *Cell*, 88:223-233) provided by the Nde1-EcoR1 fragment:

```
cat atg acc tct cgc cgc tcc gtg aag tcg ggt ccg cgg gag gtt ccg cgc gat gag tac gag gat ctg tac tac
acc ccg tct tca ggt atg gcg agt ccc gat agt ccg cct gac acc tcc cgc cgt ggc gcc cta cag aca cgc tcg
cgc cag agg ggc gag gtc cgt ttc gtc cag tac gac gag tcg gat tat gcc ctc tac ggg ggc tcg tca tcc gaa
gac gac gaa cac ccg gag gtc ccc ccg acg cgg cgt ccc gtt tcc ggg cgc gtt ttg tcc ggc ccg ggg cct gcg
cgg gcg cct ccg cca ccc gct ggg tcc gga ggg gcc gga cgc aca ccc acc acc gcc ccc cgg gcc ccc cga
acc cag cgg gtg gcg act aag gcc ccc gcg gcc ccg gcg gcg gag acc acc cgc ggc agg aaa tcg gcc cag
cca gaa tcc gcc gca ctc cca gac gcc ccc gcg tcg acg gcg cca acc cga tcc aag aca ccc gcg cag ggg
ctg gcc aga aag ctg cac ttt agc acc gcc ccc cca aac ccc gac gcg cca tgg acc ccc cgg gtg gcc ggc ttt
aac aag cgc gtc ttc tgc gcc gcg gtc ggg cgc ctg gcg gcc atg cat gcc cgg atg gcg gcg gtc cag ctc tgg
gac atg tcg cgt ccg cgc aca gac gaa gac ctc aac gaa ctc ctt ggc atc acc acc atc cgc gtg acg gtc tgc
gag ggc aaa aac ctg ctt cag cgc gcc aac gag ttg gtg aat cca gac gtg gtg cag gac gtc gac gcg gcc acg
gcg act cga ggg cgt tct gcg gcg tcg cgc ccc acc gag cga cct cga gcc cca gcc cgc tcc gat tct cgc ccc
aga cgg ccc gtc gag gaa ttc
```

[0120] which encodes the HSV-1 VP22 peptide having the sequence:

```
MTSRRSVKSGPREVPRDEYEDLYYTPSSGMA SPDSPPDTSRRGALQTRSRQRGEVRFVQ
YDESDYALYGGSSSEDEHPEVPRTRRPVSGAVLSGPGPARAPPPAGSGGAGRTPTTAP
RAPRTGRVATKAPAAPAAETTRGRKSAQPESAALPDAPASTAPTRSKTPAQGLARKLHF
STAPPNPDPAPWTPRVAGFNKRVFCAAVGRLAAMHARMAAVQLWDMSPRPTDEDLNE
LLGITTRVTVCEGKNLLQRANELVNPVDVQDVDAATATGRSAASRPTERPRAPARSA
SRPRRPVE
```

[0121] In still another embodiment, the fusion protein includes the C-terminal domain of the VP22 protein from, e.g., the nucleotide sequence (Nde1-EcoR1 fragment):

```
[0122] cat atg gac gtc gac gcg gcc acg gcg act cga ggg
cgt tct gcg gcg teg cgc ccc acc gag ega cct cga gcc cca
gcc cgc tcc gct tct cgc ccc aga egg ccc gtc gag gaa ttc
```

[0123] which encodes the VP22 (C-terminal domain) peptide sequence:

```
MDVDAATATGRSAASRPTERPRAPARSAASRPRRPVE
```

[0124] (v) Transcellular Polypeptides as General Gene Therapy Agents

[0125] Another aspect of the present invention pertains generally to the use of internalizing peptides as part of a strategy to deliver therapeutic proteins by gene therapy techniques. In addition to the subject CKI polypeptides, other therapeutic proteins such as various tumor suppressors, transcription factors, signal transduction proteins, antiviral polypeptides, and therapeutic peptides, e.g., which are otherwise localized intracellularly, can be engineered to include an internalizing peptide which drives the translocation of the therapeutic polypeptide into surrounding cells. Thus, the transcytosis system can be used to increase the efficacy of gene therapy by delivering the therapeutic protein not only to cells actually transfected with the expression vector, but also to the surrounding cells. Such constructs can be generated with other therapeutic polypeptides using such of the general teachings herein as applicable and the general knowledge in the art.

[0126] In one illustrative embodiment, the therapeutic polypeptide sequence is derived from a tumor suppressor. In addition to the CKI proteins described above, exemplary tumor suppressors from which the subject transcellular proteins can be generated include the Rb protein (and related proteins) and the p53 protein. For instance, the subject constructs can be generated with a therapeutic polypeptide domain comprising at least a functional domain of Rb, p107 or other Rb-like protein. Such tumor suppressor domains have been mapped through both genetic and biochemical means. (Hu et al. (1990) *EMBO J.* 9:1147-1155; Ewen et al. (1991) *Cell* 66:1155-1164; Ewen et al. (1992) *Science* 255:85-87). An approximately 400 amino acid fragment of Rb and p107, termed the Rb pocket, is responsible for association of these proteins with the DNA tumor virus oncoproteins and cellular ligands. Within this domain are six

regions of extensive sequence similarity between Rb and p107. (Ewen et al. (1991), *supra*).

[0127] Another class of tumor suppressors which can be used to generate the subject transcellular proteins include the DOTs (also dpc-4 and MAD-like proteins in the literature). Exemplary members of the this protein family include GenBank accession numbers U76622, U59913, U59911, U68019, U65019, U68018, U68019, 1438077, U59913 and U59912. These proteins are involved in intracellular signal transduction events downstream of TGF β (superfamily) receptors. For instance, it has been previously shown that Smad3 and Smad4 (the product of the tumor suppressor gene DPC-4) strongly synergize to induce strong ligand-independent TGF- β -like responses, and that inactive carboxy-terminally truncated mutants of either Smad act as dominant-negative inhibitors of the natural TGF- β response. Zhang et al. (1997) *Curr Biol* 7:270-276. DPC-4 has been reported as a candidate tumour suppressor. Thus, agonist and antagonist (dominant negative) forms of the DOT proteins can be used to generate transcellular polypeptides which alter the response of a cell to TGF β 's.

[0128] Another illustrative example of the subject transcellular proteins include fusion proteins generated with Bcl-2 or Bcl-x polypeptide sequences. The protein Bcl-2 plays a central role in the process of programmed cell death by blocking apoptosis. For example, when Bcl-2 levels in a cell are elevated, apoptosis is blocked. Conversely, when Bcl-2 levels in a cell are lowered, the rate of cell death is accelerated. The protein encoded by the bcl-2 proto-oncogene has been reported to be capable of inhibiting apoptosis in many hematopoietic cell systems. The bcl-2 protein is a 26 kD membrane-associated cytoplasmic protein (Tsujimoto et al. (1987) *Oncogene* 2: 3; U.S. Ser. No. 5,202,429 and U.S. Ser. No. 5,015,568; Hockenbery et al. (1991) *PNAS* 88:6961; Monaghan et al. (1992) *J. Histochem. Cytochem.* 40:1819; Nguyen et al. (1993) *J Biol. Chem.* 268: 25265; and Nguyen et al. (1994) *J. Biol. Chem.* 269:16521). The capacity of bcl-2 to enhance cell survival is related to its ability to inhibit apoptosis initiated by several factors, such as cytokine deprivation, radiation exposure, glucocorticoid treatment, and administration of anti-CD-3 antibody. Thus, all or a portion of bcl-2 sufficient to inhibit apoptosis can be used to generate the subject transcellular proteins.

[0129] Likewise, the transcellular protein can be generated using all or a portion of a protein which interacts with and/or is structurally related to the bcl-2 gene product have also been identified, such as for example bcl-x^L and bcl-x^S [Boise et al. (1993) *Cell* 74: 597; Gonzalez-Garcia et al. (1994) *Development* 120: 3033; Gottschalk et al. (1994) *PNAS* 91: 7350], Bax [Oltvai et al. (1993) *Cell* 74: 609], Mcl-1

[Kozopas et al. (1993) *PNAS* 90: 3516], and A1 [Lin et al. (1993) *J. Immunol.* 151: 179].

[0130] An example of a signal transduction protein which can be used to generate the subject transcellular proteins is the product the mammalian tubby (tub) genes, e.g., which are involved in the control of mammalian body weight. See, for example, U.S. Ser. No. 5,646,040. Tubby, an autosomal recessive mutation recently found to be the result of a splicing defect in the tubby gene. Thus, agonist and antagonist forms of tubby proteins can be used to control weight gain in animals. Moreover, these proteins are also excellent candidates for use in the treatment of ocular diseases as mutations in the tubby gene are known to lead to early progressive retinal degeneration.

[0131] Still another family of proteins which can be used to generate a transcellular protein of the present invention are the I κ B proteins. The nuclear factor- κ B (NF- κ B) is an inducible transcription factor which participates in the regulation of multiple cellular genes after treatment of cells with a variety of factors. These genes are involved in the immediate early processes of immune, acute phase, and inflammatory responses. NF- κ B has also been implicated in the transcriptional activation of several viruses, most notably the type 1 human immunodeficiency virus (HIV-1) and cytomegalovirus (CMV) (Nabel et al. (1987) *Nature* 326:711; Kaufman et al. (1987) *Mol. Cell. Biol.* 7:3759; and Sambucetti et al. (1989) *EMBO J* 8:4251).

[0132] Activation of the NF- κ B transcription factor and various related forms can be initiated by a variety of agents, including TNF- α , phorbol 12-myristate 13-acetate (PMA), interleukin-1 (IL-1) and interleukin-2 (IL-2). Activation proceeds through a post-translational event in which pre-formed cytoplasmic NF- κ B in the Rel complex is released from a cytoplasmic inhibitory protein. A common feature of the regulation of transcription factors which belong to the Rel-family is their sequestration in the cytoplasm as inactive complexes with a class of inhibitory molecules known as I κ Bs (Baeuerle et al. (1988) *Cell* 53:211; Beg et al. (1993) *Genes Dev.* 7:2064; and Gilmore et al. (1993) *Trends in Genetics* 9:427). Treatment of cells with different inducers, e.g., IL-1, TNF- α , LPS, dsRNA or PMA, results in dissociation of the cytoplasmic complexes and translocation of free NF- κ B to the nucleus (Grilli et al. (1993) *International Rev. of Cytology* 143:1-62; Baeuerle et al. (1994) *Annu. Rev. Immunol.* 12:141). The dissociation of the cytoplasmic complexes is understood to be triggered by the phosphorylation and subsequent degradation of the I κ B protein (Palombella et al. (1994) *Cell* 78:773; Ghosh et al. (1990) *Nature* 344:678).

[0133] Thus, the I κ B proteins provide a therapeutic target for being upregulated, such as by ectopic expression through gene therapy. Accordingly, the subject transcellular protein can be generated with an I κ B polypeptide sufficient to bind to and prevent nuclear localization of NF- κ B. In preferred embodiments, the I κ B polypeptide has been altered, e.g., by point mutagenesis or truncation, to increase the intracellular half-life of the fusion protein. For instance, Lys-25 and Lys-26 of human I κ B α can be mutated to remove the ability of the protein to be ubiquitinated.

[0134] In still other embodiments, the transcellular protein can be generated with a transcription factor polypeptide, as well as with a transcription repressor polypeptide, e.g., to

potentiate or inhibit the expression of a gene. In other embodiments, the therapeutic portion of the transcellular protein can be provided from a metal binding protein, e.g., for use in inhibiting the action of intracellular DNA damaging agents such as cisplatin.

[0135] In yet other embodiments, the therapeutic polypeptide of the transcellular fusion protein can be derived from small peptides/polypeptides which are artificial in sequence, or which are truncated forms of any of such proteins as described above. In an illustrative embodiment, the therapeutic polypeptide can be a peptide inhibitor of cyclin dependent kinases, such described in U.S. Ser. Pat. No. 5,625,031.

[0136] In general, the criteria for selecting the therapeutic polypeptide portion of the transcellular protein is reasonably straightforward. As with the CKI polypeptides, the therapeutic polypeptide sequence included in the fusion protein should from an intracellular protein which modulates a biological process in the cell, e.g., by either mimicking or antagonizing the wild-type form of the protein from which it is derived. The therapeutic polypeptide sequence included in the fusion protein preferably does not include any membrane association sequences, such as transmembrane domains, myristoylation sequences, etc. The therapeutic polypeptide sequence also preferably does not include any disulfide bonds. It is preferably no larger than about 100 kD, and is even more preferably no larger than 75, 50 or even 30 kD. It is also preferably no smaller than about 20 amino acids residues.

[0137] (vi) Co-delivery of CKI Polypeptides and Endothelialization Polypeptides

[0138] CKI polypeptides, including those chimeric CKI polypeptides described above, can be used to inhibit proliferation of smooth muscle cells, and can therefore be used as part of a therapeutic regimen in the treatment of a patient suffering from a condition which is characterized by excessive smooth muscle proliferation.

[0139] The arterial wall is a complex multicellular structure and is important in the regulation of inflammation, coagulation, and regional blood flow. Vascular smooth muscle cells (SMCs) are located predominantly in the arterial tunica media and are important regulators of vascular tone and blood pressure. These cells are normally maintained in a nonproliferative state in vivo. Arterial injury results in the migration of SMCs into the intimal layer of the arterial wall, where they proliferate and synthesize extracellular matrix components.

[0140] Arterial intimal thickening after injury is the result of the following series of events: 1) initiation of smooth muscle cell proliferation within hours of injury, 2) SMC migration to the intima, and 3) further SMC proliferation in the intima with deposition of matrix. Investigations of the pathogenesis of intimal thickening following arterial injury have shown that platelets, endothelial cells, macrophages and smooth muscle cells release paracrine and autocrine growth factors (such as platelet derived growth factor, PDGF α), epidermal growth factor, insulin-like growth factor, and transforming growth factor and cytokines that result in the smooth muscle cell proliferation and migration. T-cells and macrophages also migrate into the neointima. This cascade of events is not limited to arterial injury, but also

occurs following injury to veins and arterioles. The overall disease process can be termed a hyperproliferative vascular disease because of the etiology of the disease process.

[0141] Vascular injury causing intimal thickening can be broadly categorized as being either biologically or mechanically induced. One of the most commonly occurring forms of biologically mediated vascular injury leading to stenosis is Atherosclerosis. The migration and proliferation of vascular smooth muscle plays a crucial role in the pathogenesis of atherosclerosis. Atherosclerotic lesions include massive accumulation of lipid laden "foam cells" derived from monocyte/macrophage and smooth muscle cells. Formation of "foam cell" regions is associated with a breach of endothelial integrity and basal lamina destruction. Triggered by these events, restenosis is produced by a rapid and selective proliferation of vascular smooth muscle cells with increased new basal lamina (extracellular matrix) formation and results in eventual blocking of arterial pathways (Davies).

[0142] Mechanical injuries leading to intimal thickening result following balloon angioplasty, vascular surgery, transplantation surgery, and other similar invasive processes that disrupt vascular integrity. Although balloon angioplasty can dilate arterial stenosis effectively, restenosis occurs in 30-40% of patients after 6 months. Intimal thickening following balloon catheter injury has been studied in animals as a model for arterial restenosis that occurs in human patients following balloon angioplasty. De-endothelialization with an intraarterial catheter, which dilates an artery, injures the innermost layers of medial smooth muscle and may even kill some of the innermost cells.

[0143] Injury to the innermost layers of medial smooth muscle is followed by a proliferation of the medial smooth muscle cells, after which many of them migrate into the intima through fenestrae in the internal elastic lamina and proliferate to form a neointimal lesion.

[0144] Smooth muscle cell proliferation is a critical event in the pathogenesis of atherosclerosis and transplant arteriosclerosis as well as in the response to injury arising from all forms of vascular reconstruction such as angioplasty (Raines et al. (1993) *Br. Heart J.* 69 (Supplement), S 30; Clowes et al. (1991) *Vasc. Surg* 13:885; and Isik et al. (1992) *Am. J. Pathol.* 141:1139). Invasive cardiovascular surgical procedures, such as percutaneous transluminal coronary angioplasty (PTCA, using a balloon catheter) and aorto-coronary bypass surgery (ACBS), that are currently employed in treating the coronary stenosis or occlusion caused by atherosclerosis represent a major therapeutic advance for managing coronary heart disease (CHD). However, the cellular proliferative response and associated intimal hyperplasia that can follow the damage to blood vessels that occurs with these procedures leads to complications which cannot be effectively controlled by presently available drugs, and can be more detrimental than the original condition. The development of these complications, termed restenosis (in the case of PTCA) or stenosis (in the case of ACBS), has similarities to the development of atherosclerosis.

[0145] However, reendothelialization of the injured area concurrent with smooth muscle cell proliferation is a major consideration for inhibiting restenosis (Casscells, W. *Circulation* 1992, 86, 722; Reidy, M. A.; Lidner, V. in endothelial

Cell Dysfunctions, Simionescu, N. and Simionescu M., Ed. Plenum Press, NY N.Y., (1992), 31). Thus, any successful approach to inhibiting SMC proliferation should not interfere with endothelial cell repair or the normal function of other cell types (Weissberg, P. L.; Grainger, D. J.; Shanahan C. M.; Metcalfe, J. C. *Cardiovascular Res.* 1993, 27, 1191).

[0146] Accordingly, the present invention provides compositions and methods for treating patients suffering from disorders marked by unwanted proliferation of smooth muscle cells where endothelialization is a desired goal in addition to inhibiting proliferation of SMCs. In one aspect, the method of the present invention arranges for simultaneous delivery of a CKI polypeptide along with a second polypeptide (an "endothelialization polypeptide") which (a) stimulates endothelial cell proliferation; (b) stimulates migration of endothelial cells to the wound site; and/or (c) inhibits smooth muscle cell migration. For instance, in an attempt to prevent restenosis or reduced intimal smooth muscle cell proliferation following angioplasty, the present compositions can be employed clinically, concurrent with or following angioplasty.

[0147] Exemplary endothelialization polypeptides which can be co-delivered with a CKI polypeptide include angiogenic basic fibroblast growth factors (bFGF), acid fibroblast growth factor (aFGF), vascular endothelial growth factor (VEGF), platelet derived growth factor (PDGF), vascular permeability growth factor (VPF), transforming growth factor beta (TGF- β) and various cytokines. Related molecules which operate in a similar fashion may also be used. Other endothelialization polypeptides include those which may stimulate re-endothelial growth by, e.g., stimulating VEGF production by vascular smooth muscle cells, or inhibit smooth muscle cell migration by, e.g., down-regulating or antagonizing vascular smooth muscle cell PDGF receptors.

[0148] In a preferred embodiment, the endothelialization polypeptide is derived from basic FGF, such as a human bFGF. Basic FGF is an effective mitogen for vascular smooth muscle cells in the medial layer of the artery wall. For instance, a human bFGF molecule that was purified from placenta ("placental bFGF") was shown to (a) stimulate capillary endothelial cell proliferation, (b) to stimulate chemotaxis in capillary endothelial cells and (c) to stimulate these same cells to produce plasminogen activator and latent collagenase.

[0149] Another exemplary polypeptide that can be co-delivered with the subject CKIs is acidic fibroblast growth factor (aFGF). It has been demonstrated that intravenous administration of aFGF promotes vascular repair in the rat carotid artery model of restenosis, and therefore may be efficacious in the prevention of restenosis (Bjornsson et al., (1991) *PNAS* 88: 8651-8655).

[0150] In yet another preferred embodiment, the endothelialization polypeptide is derived from a vascular endothelial growth factor, particular a human VEGF.

[0151] As is clear from the context above, also included within the scope of the term "endothelialization polypeptide" are biologically active fragments thereof, as well as N-terminally or C-terminally extended versions thereof or analogs thereof substituting and/or deleting or inserting one or more amino acid residues which retain qualitatively the biological activities of the protein described herein. For

instance, preferred homologs include those in which one or more cysteine residues not required for biological activity are substituted by a different amino acid residue, preferably serine. Substitution of one or more cysteine residues reduces the opportunity for intermolecular and intramolecular disulfide bond formation, thereby rendering the molecule more stable. There are nine cysteine residues that are present in human VEGF121, bovine VEGF120, human VEGF165 and bovine VEGF164. Of these, eight are conserved with PDGF. Accordingly, the most preferred analog is one in which the ninth cysteine residue is substituted by serine. This cysteine residue is present at position 160 of hVEGF165 and position 116 of hVEGF121 and the corresponding positions in the bovine forms. Amino acid substitutions can be accomplished by site specific mutagenesis of the DNA sequences described herein using well known techniques (see, e.g., Zoller, M. J. and Smith, M., *Nucleic Acids Research* (1982) 10:6487-6500).

[0152] While the native form of the bovine vascular endothelial cell growth factor is apparently glycosylated, there is currently no evidence that glycosylation is essential for biological activity. Accordingly, biologically active non-glycosylated or partially glycosylated forms, which can be produced by prokaryotic host cells or by mutation of glycosylation sites in the protein, are included within the scope of VEGF and the other endothelialization polypeptides.

[0153] In another embodiment, the endothelialization polypeptide comprises an epidermal growth factor (EGF) polypeptide coupled with hyaluronic acid (HA). HA/EGF formulations have been demonstrated to stimulate reendothelialization in the anterior chamber of the eye. Hyaluronic acid (HA) is one of the mucopolysaccharides having a straight chain structure consisting of the repetition of a disaccharide unit of N-acetylglucosamine and glucuronic acid. HA is found in nature, in microorganisms and in the skin and connective tissue of humans and other animals.

[0154] Moreover, the glycosaminoglycans (GAGs) heparin and heparan sulfate (HS) are endogenous inhibitors of SMC proliferation, yet are able to promote endothelial cell growth (Castellot (1987) *Seminars in Thrombosis and Hemostasis* 13:489; and Wight, T. N. (1989) *Arteriosclerosis* 9:1). The endothelialization polypeptide can be derived from a protein which naturally includes HS GAG chains, or which has been engineered to artificially include HS GAG chains. See, for example, U.S. Pat. No. 5,486,599.

[0155] There are a variety of assays available for rapidly accessing the suitability of a potential endothelialization polypeptide. For instance, mitogenic activity for vascular endothelial cells can be determined by an assay which uses, as target cells, adrenal cortex-derived capillary endothelial cells (ACE cells). This assay is carried out essentially as described in Gospodarowicz et al., (1986) *J Cell Physiol.* 127:121-136). Generally, stock cultures of ACE cells are maintained in the presence of Dulbecco's modified Eagle's medium (DMEM-21) supplemented with 10% calf serum. The antibiotics penicillin (50 IU/ml), streptomycin (50 mu g/ml), gentamycin (50 mu g/ml), and Fungizone (0.25 mu g/ml) and 2 mM L-glutamine can also be added to the medium. Cells are passaged weekly on tissue culture dishes at a split ratio of between 1:40 and 1:200 (the preferred split ratio is that which gives 2.5×10^5 cells in 15 ml of medium in T75 flasks). For the mitogenic assay, cells are seeded in

12 well cluster plates at a density of 5×10^3 cells per well in 1 ml Dulbecco's modified Eagle's medium supplemented with 10% calf serum and antibiotics as described in Gospodarowicz et al., (1988) *Europ. J Cell. Biol.* 46:144-151. Alternatively, the ACE cells are plated in 35 mm dishes or 6 well cluster plates at a density of 5×10^3 cells per dish or well in 2 ml of medium as described in Gospodarowicz et al., (1986), supra. Ten-microliter aliquots of appropriate dilutions of each sample are then added to duplicate or triplicate wells in the dishes on days 0 and 2. After 4 or 5 days in culture, the plates are trypsinized and cell densities determined in a Coulter counter. For purposes of description herein, a factor is considered to have mitogenic activity for vascular endothelial cells if the cell density at the end of this assay is at least 1.5 times and preferably at least 3 times the cell density of control wells receiving no factor additions.

[0156] The CKI polypeptide and endothelialization polypeptide can be co-delivered as, for example, purified protein mixtures. The proteins can be admixed prior to administration, or administered separately but in close temporal proximity to one and other.

[0157] Alternatively, one or both of the polypeptides can be delivered by a gene therapy vector. Where both are delivered by expression vectors, the coding sequences for each polypeptide can be present on the same expression construct, or can be provided by two separate constructs, the latter requiring co-transfection with both constructs. In the instance of the former, the coding sequences for each polypeptide can be under the control of the same promoter, and may even provide for a polycistronic message (see, for example, U.S. Pat. No. 4,713,339); alternatively, the two coding sequences, while present on the same expression vector can be under the control of two separate promoters.

[0158] The CKI polypeptide can consist of as little as a CDK-binding moiety of a CKI protein, or can include a full length CKI protein and/or be a poly-CBM protein.

[0159] (vii) Exemplary Pharmaceutical Preparations

[0160] For certain of the therapeutic uses of the subject chimeric proteins, particularly cutaneous uses such as for the control of keratinocyte proliferation, direct administration of the protein will be appropriate (rather than use of a gene therapy construct). Accordingly, the subject CKI polypeptide formulations may be conveniently formulated for administration with a biologically acceptable medium, such as water, buffered saline, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like) or suitable mixtures thereof. In preferred embodiments, the chimeric protein is dispersed in lipid formulations, such as emulsions, which closely resemble the lipid composition of natural cell membranes to which the chimeric protein is to be delivered.

[0161] The optimum concentration of the active ingredient(s) in the chosen medium can be determined empirically, according to procedures well known to medicinal chemists. As used herein, "biologically acceptable medium" includes any and all solvents, dispersion media, and the like which may be appropriate for the desired route of administration of the pharmaceutical preparation. The use of such media for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the activity of the CKI polypeptide (or inter-

nalization peptide or endothelialization polypeptide as the case may be), its use in the pharmaceutical preparation of the invention is contemplated. Suitable vehicles and their formulation inclusive of other proteins are described, for example, in the book Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences. Mack Publishing Company, Easton, Pa., USA 1985).

[0162] In an exemplary embodiment, the subject CKI protein mixture is provided for transmucosal or transdermal delivery. For such administration, penetrants appropriate to the barrier to be permeated are used in the formulation with the polypeptide. Such penetrants are generally known in the art, and include, for example, for transmucosal administration bile salts and fusidic acid derivatives. In addition, detergents may be used to facilitate permeation. Transmucosal administration may be through nasal sprays or using suppositories. For topical administration, the proteins of the invention are formulated into ointments, salves, gels, or creams as generally known in the art.

[0163] In accordance with the subject method, expression constructs of the subject CKI polypeptides (and endothelialization polypeptide as appropriate) may be administered in any biologically effective carrier, e.g. any formulation or composition capable of effectively transfecting cells in vivo with a recombinant fusion gene. Approaches include insertion of the subject fusion gene in viral vectors including recombinant retroviruses, adenovirus, adeno-associated virus, and herpes simplex virus-1, or recombinant bacterial or eukaryotic plasmids. Viral vectors can be used to transfect cells directly; plasmid DNA can be delivered with the help of, for example, cationic liposomes (lipofectin) or derivatized (e.g. antibody conjugated), polylysine conjugates, gramicidin S, artificial viral envelopes or other such intracellular carriers, as well as direct injection of the gene construct or CaPO_4 precipitation carried out in vivo. It will be appreciated that because transduction of appropriate target cells represents the critical first step in gene therapy, choice of the particular gene delivery system will depend on such factors as the phenotype of the intended target and the route of administration, e.g. locally or systemically.

[0164] A preferred approach for in vivo introduction of nucleic acid encoding one of the subject CKI polypeptides into a cell is by use of a viral vector containing nucleic acid, e.g. a cDNA, encoding the gene product. Infection of cells with a viral vector has the advantage that a large proportion of the targeted cells can receive the nucleic acid. Additionally, molecules encoded within the viral vector, e.g., by a cDNA contained in the viral vector, are expressed efficiently in cells which have taken up viral vector nucleic acid.

[0165] Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery system of choice for the transfer of exogenous genes in vivo, particularly into humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. A major prerequisite for the use of retroviruses is to ensure the safety of their use, particularly with regard to the possibility of the spread of wild-type virus in the cell population. The development of specialized cell lines (termed "packaging cells") which produce only replication-defective retroviruses has increased the utility of retroviruses for gene therapy, and defective retroviruses are

well characterized for use in gene transfer for gene therapy purposes (for a review see Miller, A. D. (1990) Blood 76:271). Thus, recombinant retrovirus can be constructed in which part of the retroviral coding sequence (gag, pol, env) has been replaced by nucleic acid encoding a CKI polypeptide, rendering the retrovirus replication defective. The replication defective retrovirus is then packaged into virions which can be used to infect a target cell through the use of a helper virus by standard techniques. Protocols for producing recombinant retroviruses and for infecting cells in vitro or in vivo with such viruses can be found in *Current Protocols in Molecular Biology*, Ausubel, F. M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10-9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines for preparing both ecotropic and amphotropic retroviral systems include ψCrip , ψCre , $\psi 2$ and ψAm . Retroviruses have been used to introduce a variety of genes into many different cell types, including neural cells, epithelial cells, endothelial cells, lymphocytes, myoblasts, hepatocytes, bone marrow cells, in vitro and/or in vivo (see for example Eglitis, et al. (1985) Science 230:1395-1398; Danos and Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:6460-6464; Wilson et al. (1988) Proc. Natl. Acad. Sci. USA 85:3014-3018; Armentano et al. (1990) Proc. Natl. Acad. Sci. USA 87:6141-6145; Huber et al. (1991) Proc. Natl. Acad. Sci. USA 88:8039-8043; Ferry et al. (1991) Proc. Natl. Acad. Sci. USA 88:8377-8381; Chowdhury et al. (1991) Science 254:1802-1805; van Beusechem et al. (1992) Proc. Natl. Acad. Sci. USA 89:7640-7644; Kay et al. (1992) Human Gene Therapy 3:641-647; Dai et al. (1992) Proc. Natl. Acad. Sci. USA 89:10892-10895; Hwu et al. (1993) J. Immunol. 150:4104-4115; U.S. Pat. Nos. 4,868,116; 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573).

[0166] In choosing retroviral vectors as a gene delivery system for the subject fusion proteins, it is important to note that a prerequisite for the successful infection of target cells by most retroviruses, and therefore of stable introduction of the recombinant gene, is that the target cells must be dividing. In general, this requirement will not be a hindrance to use of retroviral vectors to deliver the subject fusion gene constructs. In fact, such limitation on infection can be beneficial in circumstances where the tissue (e.g. nontransformed cells) surrounding the target cells does not undergo extensive cell division and is therefore refractory to infection with retroviral vectors.

[0167] Furthermore, it has been shown that it is possible to limit the infection spectrum of retroviruses and consequently of retroviral-based vectors, by modifying the viral packaging proteins on the surface of the viral particle (see, for example PCT publications WO93/25234, WO94/06920, and WO94/11524). For instance, strategies for the modification of the infection spectrum of retroviral vectors include: coupling antibodies specific for cell surface antigens to the viral env protein (Roux et al. (1989) PNAS 86:9079-9083; Julian et al. (1992) J. Gen Virol 73:3251-3255; and Goud et al. (1983) Virology 163:251-254); or coupling cell surface ligands to the viral env proteins (Neda et al. (1991) J Biol Chem 266:14143-14146). Coupling can be in the form of the chemical cross-linking with a protein or other variety (e.g. lactose to convert the env protein to an asialoglycoprotein),

as well as by generating fusion proteins (e.g. single-chain antibody/env fusion proteins). This technique, while useful to limit or otherwise direct the infection to certain tissue types, and can also be used to convert an ecotropic vector in to an amphotropic vector.

[0168] Moreover, use of retroviral gene delivery can be further enhanced by the use of tissue- or cell-specific transcriptional regulatory sequences which control expression of the fusion gene of the retroviral vector.

[0169] Another viral gene delivery system useful in the present invention utilizes adenovirus-derived vectors. The genome of an adenovirus can be manipulated such that it encodes a gene product of interest, but is inactivated in terms of its ability to replicate in a normal lytic viral life cycle (see, for example, Berkner et al. (1988) *BioTechniques* 6:616; Rosenfeld et al. (1991) *Science* 252:431-434; and Rosenfeld et al. (1992) *Cell* 68:143-155). Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (e.g., Ad2, Ad3, Ad7 etc.) are well known to those skilled in the art. Recombinant adenoviruses can be advantageous in certain circumstances in that they are not capable of infecting nondividing cells and can be used to infect a wide variety of cell types, including endothelial cells (Lemarchand et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:6482-6486), and smooth muscle cells (Quantin et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584). Furthermore, the virus particle is relatively stable and amenable to purification and concentration, and as above, can be modified so as to affect the spectrum of infectivity. Additionally, introduced adenoviral DNA (and foreign DNA contained therein) is not integrated into the genome of a host cell but remains episomal, thereby avoiding potential problems that can occur as a result of insertional mutagenesis in situations where introduced DNA becomes integrated into the host genome (e.g., retroviral DNA). Moreover, the carrying capacity of the adenoviral genome for foreign DNA is large (up to 8 kilobases) relative to other gene delivery vectors (Berkner et al., supra; Haj-Ahmand and Graham (1986) *J. Virol.* 57:267). Most replication-defective adenoviral vectors currently in use and therefore favored by the present invention are deleted for all or parts of the viral E1 and E3 genes but retain as much as 80% of the adenoviral genetic material (see, e.g., Jones et al. (1979) *Cell* 16:683; Berkner et al., supra; and Graham et al. in *Methods in Molecular Biology*, E. J. Murray, Ed. (Humana, Clifton, N.J., 1991) vol. 7, pp. 109-127). Expression of the inserted fusion gene can be under control of, for example, the E1A promoter, the major late promoter (MLP) and associated leader sequences, the E3 promoter, or exogenously added promoter sequences.

[0170] Yet another viral vector system useful for delivery of the subject CKI polypeptides is the adeno-associated virus (AAV). Adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle. (For a review see Muzyczka et al. *Curr. Topics in Micro. and Immunol.* (1992) 158:97-129). It is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration (see for example Flotte et al. (1992) *Am. J. Respir. Cell. Mol. Biol.* 7:349-356; Samulski et al. (1989) *J. Virol.* 63:3822-3828; and McLaughlin et al. (1989) *J. Virol.* 62:1963-1973). Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate.

Space for exogenous DNA is limited to about 4.5 kb. An AAV vector such as that described in Tratschin et al. (1985) *Mol. Cell. Biol.* 5:3251-3260 can be used to introduce DNA into cells. A variety of nucleic acids have been introduced into different cell types using AAV vectors (see for example Hermonat et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6466-6470; Tratschin et al. (1985) *Mol. Cell. Biol.* 4:2072-2081; Wondisford et al. (1988) *Mol. Endocrinol.* 2:32-39; Tratschin et al. (1984) *J. Virol.* 51:611-619; and Flotte et al. (1993) *J. Biol. Chem.* 268:3781-3790).

[0171] Other viral vector systems that may have application in gene therapy have been derived from herpes virus, vaccinia virus, and several RNA viruses. In particular, herpes virus vectors may provide a unique strategy for persistent expression of the subject fusion proteins in cells of the central nervous system and ocular tissue (Pepose et al. (1994) *Invest Ophthalmol Vis Sci* 35:2662-2666).

[0172] In addition to viral transfer methods, such as those illustrated above, non-viral methods can also be employed to cause expression of a the subject proteins in the tissue of an animal. Most nonviral methods of gene transfer rely on normal mechanisms used by mammalian cells for the uptake and intracellular transport of macromolecules. In preferred embodiments, non-viral gene delivery systems of the present invention rely on endocytic pathways for the uptake of the gene by the targeted cell. Exemplary gene delivery systems of this type include liposomal derived systems, poly-lysine conjugates, and artificial viral envelopes.

[0173] In a representative embodiment, a gene encoding one of the subject proteins can be entrapped in liposomes bearing positive charges on their surface (e.g., lipofectins) and (optionally) which are tagged with antibodies against cell surface antigens of the target tissue (Mizuno et al. (1992) *No Shinkei Geka* 20:547-551; PCT publication WO91/06309; Japanese patent application 1047381; and European patent publication EP-A-43075). For example, lipofection of neuroglioma cells can be carried out using liposomes tagged with monoclonal antibodies against glioma-associated antigen (Mizuno et al. (1992) *Neurol. Med. Chir.* 32:873-876).

[0174] In yet another illustrative embodiment, the gene delivery system comprises an antibody or cell surface ligand which is cross-linked with a gene binding agent such as poly-lysine (see, for example, PCT publications WO93/04701, WO92/22635, WO92/20316, WO92/19749, and WO92/06180). For example, the subject gene construct can be used to transfect hepatocytic cells in vivo using a soluble polynucleotide carrier comprising an asialoglycoprotein conjugated to a polycation, e.g. poly-lysine (see U.S. Pat. No. 5,166,320). It will also be appreciated that effective delivery of the subject nucleic acid constructs via receptor-mediated endocytosis can be improved using agents which enhance escape of the gene from the endosomal structures. For instance, whole adenovirus or fusogenic peptides of the influenza HA gene product can be used as part of the delivery system to induce efficient disruption of DNA-containing endosomes (Mulligan et al. (1993) *Science* 260:926; Wagner et al. (1992) *PNAS* 89:7934; and Christiano et al. (1993) *PNAS* 90:2122).

[0175] In clinical settings, the gene delivery systems can be introduced into a patient by any of a number of methods, each of which is familiar in the art. For instance, a pharma-

ceutical preparation of the gene delivery system can be introduced systemically, e.g. by intravenous injection, and specific transduction of the target cells occurs predominantly from specificity of transfection provided by the gene delivery vehicle, cell-type or tissue-type expression due to the transcriptional regulatory sequences controlling expression of the gene, or a combination thereof. In other embodiments, initial delivery of the recombinant gene is more limited with introduction into the animal being quite localized. For example, the gene delivery vehicle can be introduced by catheter (see U.S. Pat. No. 5,328,470) or by stereotactic injection (e.g. Chen et al. (1994) PNAS 91: 3054-3057).

[0176] Moreover, the pharmaceutical preparation can consist essentially of the gene delivery system in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery system can be produced in tact from recombinant cells, e.g. retroviral packages, the pharmaceutical preparation can comprise one or more cells which produce the gene delivery system. In the case of the latter, methods of introducing the viral packaging cells may be provided by, for example, rechargeable or biodegradable devices. Various slow release polymeric devices have been developed and tested in vivo in recent years for the controlled delivery of drugs, including proteinacious biopharmaceuticals, and can be adapted for release of viral particles through the manipulation of the polymer composition and form. A variety of biocompatible polymers (including hydrogels), including both biodegradable and non-degradable polymers, can be used to form an implant for the sustained release of an the viral particles by cells implanted at a particular target site. Such embodiments of the present invention can be used for the delivery of an exogenously purified virus, which has been incorporated in the polymeric device, or for the delivery of viral particles produced by a cell encapsulated in the polymeric device.

[0177] By choice of monomer composition or polymerization technique, the amount of water, porosity and consequent permeability characteristics can be controlled. The selection of the shape, size, polymer, and method for implantation can be determined on an individual basis according to the disorder to be treated and the individual patient response. The generation of such implants is generally known in the art. See, for example, Concise Encyclopedia of Medical & Dental Materials, ed. by David Williams (MIT Press: Cambridge, Mass., 1990); and the Sabel et al. U.S. Pat. No. 4,883,666. In another embodiment of an implant, a source of cells producing the recombinant virus is encapsulated in implantable hollow fibers. Such fibers can be pre-spun and subsequently loaded with the viral source (Aebischer et al. U.S. Pat. No. 4,892,538; Aebischer et al. U.S. Pat. No. 5,106,627; Hoffman et al. (1990) Expt. Neurobiol. 110:39-44; Jaeger et al. (1990) Prog. Brain Res. 82:41-46; and Aebischer et al. (1991) J. Biomech. Eng. 113:178-183), or can be co-extruded with a polymer which acts to form a polymeric coat about the viral packaging cells (Lim U.S. Pat. No. 4,391,909; Sefton U.S. Pat. No. 4,353,888; Sugamori et al. (1989) Trans. Am. Artif. Intern. Organs 35:791-799; Sefton et al. (1987) Biotechnol. Bioeng. 29:1135-1143; and Aebischer et al. (1991) Biomaterials 12:50-55). Again, manipulation of the polymer can be carried out to provide for optimal release of viral particles.

[0178] (viii) Exemplary Uses for the Subject Antiproliferative Formulations

[0179] The subject fusion proteins can be used in the treatment of hyperproliferative vascular disorders, e.g. smooth muscle hyperplasia (such as atherosclerosis) or restinosis, as well as other disorders characterized by fibrosis, e.g. rheumatoid arthritis, insulin dependent diabetes mellitus, glomerulonephritis, cirrhosis, and scleroderma, particularly proliferative disorders in which loss of TGF- β autocrine or paracrine signaling, and accordingly loss of p15 function, is implicated.

[0180] For example, restinosis continues to limit the efficacy of coronary angioplasty despite various mechanical and pharmaceutical interventions that have been employed. An important mechanism involved in normal control of intimal proliferation of smooth muscle cells appears to be the induction of autocrine and paracrine TGF- β inhibitory loops in the smooth muscle cells (Scott-Burden et al. (1994) Tex Heart Inst J 21:91-97; Grainger et al. (1993) Cardiovasc Res 27:2238-2247; and Grainger et al. (1993) Biochem J 294:109-112). Loss of sensitivity to TGF- β , or alternatively, the overriding of this inhibitory stimulus such as by PDGF autostimulation, can be a contributory factor to abnormal smooth muscle proliferation in restinosis. It may therefore be possible to treat or prevent restinosis by the use of gene therapy with CDK inhibitor fusion protein of the present invention. The fusion gene construct can be delivered, for example, by percutaneous transluminal gene transfer (Mazur et al. (1994) Tex Heart Inst J 21:104-111) using viral or liposomal delivery compositions. An exemplary adenovirus-mediated gene transfer technique and compositions for treatment of cardiac or vascular smooth muscle is provided in PCT publication WO 94/11506.

[0181] In one embodiment, the co-delivery of a CKI polypeptide with an endothelialization polypeptide can be used in the treatment of disorders/injury marked by unwanted smooth muscle cell proliferation and deepithelialization. For example, the co-administration approach can be used as a post-operative wound healing agent in balloon angioplasty, a procedure in which vascular endothelial cells are removed or damaged, together with compression of atherosclerotic plaques. The two polypeptides can be co-delivered to inner vascular surfaces by systemic or local intravenous application either as intravenous bolus injection or infusions of polypeptides, expression constructs, or a mixture thereof. If desired, the endothelialization polypeptide can be administered over time using a micrometering pump, while the CKI polypeptide is ultimately delivered by gene therapy. Suitable compositions for intravenous administration comprise the endothelialization polypeptide in an amount effective to promote endothelial cell growth and a parenteral carrier material. The endothelialization polypeptide can be present in the composition over a wide range of concentration, administered once or in dosing regimens that allow for multiple applications. Any of the known parenteral carrier vehicles can be used, such as normal saline or 5-10% dextrose.

[0182] The co-delivery embodiment of the subject method can also be used to inhibit SMC proliferation yet promote endothelialization in vascular graft surgery. In the case of vascular grafts using either transplanted vessels or synthetic material, for example, the endothelialization polypeptide can be applied to the surfaces of the graft and/or at the junctions

of the graft and the existing vasculature in order to promote the growth of vascular endothelial cells. For such applications, the endothelialization polypeptide can be applied intravenously as described above for balloon angioplasty or it can be applied directly to the surfaces of the graft and/or the existing vasculature either before or during surgery.

[0183] Transforming growth factor- β is also understood to play a significant role in local glomerular and interstitial sites in human kidney development and disease. Consequently, the subject method provides a method of treating or inhibiting glomerulopathies and other renal proliferative disorders comprising the in vivo delivery and recombinant expression of the subject fusion proteins, e.g., the poly-CBM and/or transcellular CKI polypeptides, in kidney tissue.

[0184] In other embodiments, therapeutic application of a CDK inhibitor fusion protein, e.g., by gene therapy, can be used in the treatment of a neuroglioma. Gliomas account for 40-50% of intracranial tumors at all ages of life. Despite the increasing use of radiotherapy, chemotherapy, and sometimes immunotherapy after surgery for malignant glioma, the mortality and morbidity rates have not substantially improved. However, there is increasing experimental and clinical evidence that for a significant number of gliomas, loss of TGF- β responsiveness is an important event in the loss of growth control. Irrespective of the cause of decreased responsiveness, e.g. the loss of function of p15 or the loss of other TGF- β signal transduction proteins, exogenous expression of, for example, an INK4 fusion protein such as p15/p27 fusion protein in the cell can be used effectively to inhibit cell proliferation.

[0185] Yet another aspect of the invention pertains to methods of treating proliferative and/or differentiative disorders which arise from cells which, despite aberrant growth control, still require one or more CDKs (e.g., CDK4 or CDK6) for cell growth. There are a wide variety of pathological cell proliferative conditions for which the fusion gene constructs of the present invention can provide therapeutic benefits, with the general strategy being the inhibition of an anomalous cell proliferation. For instance, the gene constructs of the present invention can be used as a part of a gene therapy protocol in a cell in which a cell-cycle regulatory protein (such as an INK4 or CIP protein) is misexpressed or in which signal transduction pathways upstream of the protein are dysfunctional. To illustrate, cell types which exhibit pathological or abnormal growth presumably dependent at least in part on a function of a, INK4 or CIP protein include various cancers and leukemias, psoriasis, bone diseases, fibroproliferative disorders such as involving connective tissues, atherosclerosis and other smooth muscle proliferative disorders, as well as chronic inflammation. In addition to proliferative disorders, the treatment of differentiative disorders which result from, for example, de-differentiation of tissue which may (optionally) be accompanied by abortive reentry into mitosis. Such degenerative disorders include chronic neurodegenerative diseases of the nervous system, including Alzheimer's disease, Parkinson's disease, Huntington's chorea, amyotrophic lateral sclerosis and the like, as well as spinocerebellar degenerations. Other differentiative disorders include, for example, disorders associated with connective tissue, such as may occur due to de-differentiation of chondrocytes or osteocytes, as well as vascular disorders which involve de-differentiation of endothelial tissue and smooth

muscle cells, gastric ulcers characterized by degenerative changes in glandular cells, and renal conditions marked by failure to differentiate, e.g. Wilm's tumors. It will also be apparent that, by transient use of gene therapy constructs of the subject fusion proteins, in vivo reformation of tissue can be accomplished, e.g. in the development and maintenance of organs. By controlling the proliferative and differentiative potential for different cells, the subject gene constructs can be used to reform injured tissue, or to improve grafting and morphology of transplanted tissue. For example, the subject CDK inhibitors can be employed therapeutically as part of a regimen to regulate organs after physical, chemical or pathological insult.

[0186] Furthermore, as described in the art, transformation of a cell can be due in part to a loss-of-function mutation to a particular INK4 gene, e.g., ranging from a point mutation to gross deletion of the gene. Additionally, other data suggests that certain disorders may arise because cells have lost the ability to induce expression of an INK4 gene. Normal cell proliferation, for instance, is generally marked by responsiveness to negative autocrine or paracrine growth regulators, such as members of the TGF- β family, e.g. TGF- β 1, TGF- β 2 or TGF- β 3, and related polypeptide growth inhibitors. Ordinarily, control of cellular proliferation by such growth regulators, particularly in epithelial and hemopoietic cells, is in the form of growth inhibition. Moreover, as described in Hannon and Beach (1995) *Nature* 371:257-261, TGF- β inhibits cell proliferation by inducing expressions of p15, which in turn inhibits activation of CDK4 or CDK6 complexes.

[0187] It has been observed that a significant percentage of human cancers derived from cells types ordinarily inhibited by TGF- β display a reduced responsiveness to this growth regulator. For instance, some tumors of colorectal, liver epithelial, and epidermal origin show reduced sensitivity and resistance to the growth-inhibitory effects of TGF- β as compared to their normal counterparts. In this context, a noteworthy characteristic of several retinoblastoma cell lines is the absence of detectable TGF- β receptors. Treatment of such tumors with the subject fusion proteins provides an opportunity to mimic the TGF- β inhibitory signal. Moreover, it will be appreciated that the subject method can be used generally to inhibit proliferation of cells which, in general, are still reliant on cyclin dependent kinases.

[0188] It has been demonstrated that gene therapy can be used to target glioma cells for expression of recombinant proteins (Miyao et al. (1993) *J. Neurosci. Res.* 36:472-479; Chen et al. (1994) *PNAS* 91:3054-3057; and Takamiya et al. (1993) *J. Neurosurg.* 79:104-110). Thus, a gene construct for expressing the subject fusion protein can be delivered to the tumor, preferably by stereotactic-dependent means. In preferred embodiments, the gene delivery system is a retroviral vector. Since rapidly growing normal cells are rare in the adult CNS, glioma cells can be specifically transduced with a recombinant retrovirus. For example, the retroviral particle can be delivered into the tumor cavity through an Ommaya tube after surgery, or alternatively, packaging fibroblasts encapsulated in retrievable immunoisulatory vehicles can be introduced into the tumor cavity. In order to increase the effectiveness and decrease the side effects of the retrovirus-mediated gene therapy, glioma-specific promoters can be used to regulate expression of the therapeutic gene. For example, the promoter regions of glial fibrillary acidic

protein (GFAP) and myelin basis protein (MBP) can operably linked to the fusion gene in order to direct glial cell-specific expression of the fusion protein.

[0189] In another embodiment, gene therapy can be used in conjunction with the subject fusion proteins in the treatment of various carcinomas. In a representative embodiment, a gene therapy system comprising the subject fusion gene is used to treat certain breast cancers. In preferred embodiments, expression of the subject fusion protein is controlled at least in part by a mammary-specific promoter, a number of which are available (for review, see Hennighausen (1990) *Protein Expression and Purification* 1:3-8; and Günzberg et al. (1992) *Biochem J* 283:625-632).

[0190] In similar fashion, gene therapy protocols involving delivery of the subject fusion protein can be used in the treatment of malignant melanoma, which also serves as a model for progressive TGF- β resistance in transformation. In preferred embodiments, gene therapy protocols for treatment of melanomas include, in addition to the delivery of the fusion gene construct, the delivery of a pharmaceutical preparation of the gene by direct injection. For instance, U.S. Pat. No. 5,318,514 describes an applicator for the electroporation of genes into epidermal cells and can be used in accordance with the present invention.

[0191] The subject method can also be used to treat retinoblastomas in which the retinoblastoma gene (RB) is not itself impaired, e.g. the effective impairment of the RB checkpoint is the result of a failure to control CDK4 phosphorylation of RB. Thus, one of the subject fusion proteins can be expressed in a retinoblastoma cell, thereby causing inhibition of CDK4 activation and down-regulating RB phosphorylation. To illustrate, a recombinant retrovirus can be constructed to facilitate expression of a fusion protein including an INK4 protein, e.g., derived from p16 or p15, and a CIP protein, e.g., derived from p21, p27 or p57. Infectivity of retinoblastoma cells can be enhanced by derivatizing the env protein with antibodies specific for retinoblastoma cells, e.g. antibodies to retinal S-antigen (Doroso et al. (1985) *Invest Ophthalmol Vis Sci* 26:560-572; see also Liao et al. (1981) *Eur J Immunol* 11:450-454; and U.S. Pat. No. 4,444,744).

[0192] In yet another embodiment, the subject gene is delivered to a sarcoma, e.g. an osteosarcoma or Kaposi's sarcoma. In a representative embodiment, the gene is provided in a viral vector and delivered by way of a viral particle which has been derivatized with antibodies immunoselective for an osteosarcoma cell (see, for example, U.S. Pat. Nos. 4,564,517 and 4,444,744; and Singh et al. (1976) *Cancer Res* 36:4130-4136).

[0193] Given the role of CDK activation in various epithelial cell proliferative disorders, it will be evident that the subject fusion proteins will find ready application for the treatment or prophylaxis of, for example, psoriasis; keratosis; acne; comedogenic lesions; verrucous lesions such as verruca plana, plantar warts, verruca acuminata, and other verruciform lesions marked by proliferation of epithelial cells; folliculitis and pseudofolliculitis; keratoacanthoma; callosities; Darier's disease; ichthyosis; lichen planus; molluscous contagiosum; melasma; Fordyce disease; and keloids or hypertrophic scars.

[0194] Yet another aspect of the present invention relates to the use of the subject fusion proteins to control hair

growth. The growth of hard keratin fibers such as wool and hair is dependent on the proliferation of dermal sheath cells. Hair follicle stem cells of the sheath are highly active, and give rise to hair fibers through rapid proliferation and complex differentiation. The hair cycle involves three distinct phases: anagen (growing), catagen (regressing), and telogen (resting). The epidermal stem cells of the hair follicle are activated by dermal papilla during late telogen. This is termed "bulge activation". Moreover, such stem cells are thought to be pluripotent stem cells, giving rise not only to hair and hair follicle structures, but also the sebaceous gland and epidermis. The subject method provides a means for altering the dynamics of the hair growth cycle to induce quiescence of proliferation of hair follicle cells, particularly stem cells of the hair follicle, inhibiting CDK activation.

[0195] For instance, gene therapy treatments or, alternatively, topical administration of a fusion protein preparation, can be employed as a way of reducing the growth of human hair as opposed to its conventional removal by cutting, shaving, or depilation. For instance, the present method can be used in the treatment of trichosis characterized by abnormally rapid or dense growth of hair, e.g. hypertrichosis. In an exemplary embodiment, the subject fusion proteins can be used to manage hirsutism, a disorder marked by abnormal hairiness. Application of the CDK inhibitors of the present invention can also provide a process for extending the duration of depilation.

[0196] Moreover, because the CDK inhibitor fusion proteins are likely to be cytostatic to epithelial cells, rather than cytotoxic, these proteins can be used to protect hair follicle cells from cytotoxic agents which require progression into S-phase of the cell-cycle for efficacy, e.g. radiation-induced death. Treatment with a CDK inhibitor of the present invention provides protection by causing the hair follicle cells to become quiescent, e.g., by inhibiting the cells from entering S phase, and thereby preventing the follicle cells from undergoing mitotic catastrophe or programmed cell death. For instance, such treatments can be used for patients undergoing chemo- or radiation-therapies which ordinarily result in hair loss.

[0197] The subject method can also be used in the treatment of folliculitis, such as folliculitis decalvans, folliculitis ulerythematosus reticulata or keloid folliculitis. For example, a cosmetic preparation of an CDK inhibitory fusion protein can be applied topically in the treatment of pseudofolliculitis, a chronic disorder occurring most often in the submandibular region of the neck and associated with shaving, the characteristic lesions of which are erythematous papules and pustules containing buried hairs.

[0198] In similar fashion, such preparations can be used in the treatment of granulomas, e.g. tumor-like mass or nodule of granulation tissue, which may include epithelial tissue derived from cutaneous or mucosal sources.

[0199] In another aspect of the invention, the subject method can be used in conjunction with various periodontal procedures in which inhibition of epithelial cell proliferation in and around periodontal tissue is desired. For example, preparations of the present invention can find application in the treatment of periodontal disease. It is estimated that in the United States alone, there are in excess of 125 million adults with periodontal disease in varying forms. Periodontal disease starts as inflammatory lesions because of specific

bacteria localizing in the area where the gingiva attaches to the tooth. Usually first to occur is a vascular change in the underlying connective tissue. Inflammation in the connective tissue stimulates the following changes in the epithelial lining of the sulcus and in the epithelial attachment: increased mitotic activity in the basal epithelial layer; increased producing of keratin with desquamation; cellular desquamation adjacent to the tooth surface tends to deepen the pocket; epithelial cells of the basal layer at the bottom of the sulcus and in the area of attachment proliferate into the connective tissue and break up of the gingival fibers begins to occur, wherein dissolution of the connective tissue results in the formation of an open lesion. The application of CDK inhibitor preparations to the periodontium can be used to inhibit proliferation of epithelial tissue and thus prevent further periodontoclastic development.

[0200] In yet another embodiment of the present invention, the subject CDK inhibitors can be used to inhibit spermatogenesis or oogenesis by inhibiting progression through mitotic or meiotic cell-cycle stages. The anti-mitotic and/or anti-meiotic activity of the fusion proteins identified in the present invention may accordingly be used, for example, in birth control methods by disrupting oogenic pathways in order to prevent the development of either the egg or sperm, or by preventing mitotic progression of a fertilized egg.

[0201] In a still further embodiment, the subject fusion protein is recombinantly expressed in tissue which is characterized by unwanted de-differentiation and which may also be undergoing unwanted apoptosis. For instance, many neurological disorders are associated with degeneration of discrete populations of neuronal elements. For example, Alzheimer's disease is associated with deficits in several neurotransmitter systems, both those that project to the neocortex and those that reside with the cortex. For instance, the nucleus basalis in patients with Alzheimer's disease were observed to have a profound (75%) loss of neurons compared to age-matched controls. Although Alzheimer's disease is by far the most common form of dementia, several other disorders can produce dementia. Many are age-related, occurring in far greater incidence in older people than in younger. Several of these are degenerative diseases characterized by the death of neurons in various parts of the central nervous system, especially the cerebral cortex. However, some forms of dementia are associated with degeneration of the thalamus or the white matter underlying the cerebral cortex. Here, the cognitive dysfunction results from the isolation of cortical areas by the degeneration of efferents and afferents. Huntington's disease involves the degeneration of intrastriatal and cortical cholinergic neurons and GABAergic neurons. Pick's disease is a severe neuronal degeneration in the neocortex of the frontal and anterior temporal lobes, sometimes accompanied by death of neurons in the striatum. Accordingly, the subject fusion proteins can be delivered to the effected tissue by gene therapy techniques. It is noted that numerous advances have been made in the construction of expression vectors, cellular and viral transgene carriers, and the characterization of target cells for neuronal gene therapy, and can be readily adapted for delivery of the subject genes (see, for example, Suhr et al. (1993) Arch Neurol 50:1252-1268; Jiao et al. (1993) Nature 362:450-453; Friedmann (1992) Ann Med 24:411-417; and Freese et al. (1991) Nuc Acid Res 19:7219-7223).

[0202] In addition to degenerative-induced dementias, the subject gene therapy systems can be applied opportunely in the treatment of neurodegenerative disorders which have manifestations of tremors and involuntary movements. Parkinson's disease, for example, primarily affects subcortical structures and is characterized by degeneration of the nigrostriatal pathway, raphe nuclei, locus cereleus, and the motor nucleus of vagus. Ballism is typically associated with damage to the subthalamic nucleus, often due to acute vascular accident. Also included are neurogenic and myopathic diseases which ultimately affect the somatic division of the peripheral nervous system and are manifest as neuromuscular disorders. Examples include chronic atrophies such as amyotrophic lateral sclerosis, Guillain-Barre syndrome and chronic peripheral neuropathy, as well as other diseases which can be manifest as progressive bulbar palsies or spinal muscular atrophies. Moreover, the use of the subject fusion gene therapy constructs is amenable to the treatment of disorders of the cerebellum which result in hypotonia or ataxia, such as those lesions in the cerebellum which produce disorders in the limbs ipsilateral to the lesion. For instance, p16/p27 fusion gene constructs can be used to treat a restricted form of cerebellar cortical degeneration involving the anterior lobes (vermis and leg areas) such as is common in alcoholic patients.

[0203] Furthermore, the subject fusion proteins can also be used in the treatment of autonomic disorders of the peripheral nervous system, which include disorders affecting the innervation of smooth muscle and endocrine tissue (such as glandular tissue). For instance, recombinant fusion protein of the present invention can be expressed by gene therapy and used to treat tachycardia or atrial cardiac arrhythmias which may arise from a degenerative condition of the nerves innervating the striated muscle of the heart.

[0204] As will be apparent, the subject gene constructs can be used to cause expression of the fusion polypeptides in cells propagated in culture, e.g. to produce proteins or polypeptides, including fusion proteins or polypeptides, for purification. In addition, recombinant expression of the subject fusion polypeptides in cultured cells can be useful for controlling differentiation states of cells in vitro, for instance, by controlling the level of activation of a CDK. To illustrate, in vitro neuronal culture systems have proved to be fundamental and indispensable tools for the study of neural development, as well as the identification of neurotrophic factors. Once a neuronal cell has become terminally-differentiated, it typically will not change to another terminally differentiated cell-type. However, neuronal cells can nevertheless readily lose their differentiated state. This is commonly observed when they are grown in culture from adult tissue, and when they form a blastema during regeneration. By preventing the activation of one or more CDKs, particularly in G₀ or G₁, certain of the subject fusion proteins can prevent mitotic progression and hence provide a means for ensuring an adequately restrictive environment in order to maintain neuronal cells at various stages of differentiation, and can be employed, for instance, in cell cultures designed to test the specific activities of trophic factors. Other tissue culture systems which require maintenance of differentiation will be readily apparent to those skilled in the art. In this respect, each of the subject antagonist of CDK4 activation can be used for ex vivo tissue generation, as for example, to enhance the generation of prosthetic tissue devices for implantation. That is, by inhibiting the activation

of a CDK with one of the subject fusion proteins, cultured cells can be guided along certain differentiative pathways.

EXEMPLIFICATION

[0205] The invention now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

[0206] Prototype embodiments of the CDK inhibitory fusion protein described above were derived from the fusion of the coding sequences from the human p27 and p16 cDNAs. The nucleotide sequence for the fusion gene encoding the p27-p16 protein is provided in SEQ ID No. 1, with the corresponding amino acid sequence being designated by SEQ ID No. 2. The construct includes a poly(His) leader for purification, along with a hinge region including a (Gly,Ser)₃ linker to permit proper folding and breathing of each of the p27 and p16 portions of the resulting protein. The sequences for both human p27 and human p16 have been described in the art. Briefly, the p27-p16 fusion protein was constructed as follow.

[0207] The expression vector is pT7-7 from US Biochemical. To construct the p27-p16 fusion, first we PCR amplified the p27 coding sequence using the following primers:

[0208] N-terminal primer: (SEQ ID No. 3)

5'-GCGGCCGGTCATATGCACCACCATCACCATCAACG-TGCGAGTGTCT-3'

[0209] This primer carries an NdeI site and 6 histidine codons that are inserted between the ATG and the second amino acid of p27.

[0210] C-terminal primer: (SEQ ID No. 4)

5'-GCCGCCGGCGTCGACTCGCCGAATTCGGATCCACCCCGCCGAA-
CGCCACCCCGCGTGCCTCCGCCACCGTTGACGTCTCTTGAGGCCAG-
G-3'

[0211] This primer carries the (Gly,Ser)₃ repeat and EcoRI, SalI and Hind3 restriction sites and eliminates the stop codon of p27.

[0212] The p27 PCR product was cut with NdeI and Hind3 and inserted into pT7-7 cut with NdeI and Hind3. The resulted construct was cut with EcoRI and SalI and a full length p16 PCR product was inserted as an EcoRI-XhoI fragment. The position of the EcoRI site allows the in-frame insertion of p16. The rest of the hinge region between the p27 and p16 coding sequences derives from the 5' end of the p16 cDNA.

[0213] The pT7p27-p16 expression plasmid was transformed into BL21 cells. For fusion protein expression, cells were grown in LB+50 µg/ml ampicillin at 37 C. to OD₆₀₀=0.8 and protein expression was induced by IPTG (final; conc.: 20 mM) for 4 hours at 37 C. Cells were collected and the pellet was frozen at -80 C. The preparation of the cell lysate and binding to a Ni²⁺ charged sepharose resin (Invit-

rogen catalog no. R801) was done according to the manufacturer's instruction (Invitrogen; see also Hochuli et al. (1987) J. Chromatography 411:177-184; and Janknecht et al. (1991) PNAS 88:8972-8976). The bound proteins were eluted with 50 mM, 200 mM, 350 mM, and 500 mM imidazol and the fractions were analyzed on SDS/PAGE. The 200 mM, 350 mM, and 500 mM imidazol fractions were collected, dialysed against 1xPBS(1 mM KH₂PO₄, 10 mM Na₂HPO₄, 137 mM NaCl, 2.7 mM KCl, pH=7.4)+10% glycerol and stored at -80 C. in aliquots. ~25% of the prep was the fusion protein.

[0214] The purity of the p27-p16, p27, and p16 preparations were normalized using p16 and p27 specific antibodies.

[0215] The kinase inhibitory activity of the p27-p16 fusion protein was determined using an in vitro kinase assay in which the kinase activity of a particular cyclin/CDK complex was measured for varying concentrations of fusion protein. Briefly, the assay employs Sf9 cell extracts that were made from cells that were coinfectd with the proper CDK and cyclin expression constructs. Typically, 44 µg of Sf9 extract in 50 µl of 50 mM Tris/Cl pH=7.6, 10 mM MgCl₂, 1 mM DTT, 25 µM ATP, 10 µCi ³²P-γ-ATP was used in the absence of the presence of the particular inhibitor (inhibitor concentration was between 25 nM to 1 µM). The reaction was carried out at 30° C. for 30 minutes using 2 µg of Gst-Rb as a substrate. Gst-Rb was recaptured using GSH-agarose, separated on 10% SDS/PAGE and stained with Comassie

blue. After autoradiography the GST-Rb bands were cut out and ³²P incorporation was measured.

[0216] The concentration of p27-p16 fusion protein at which 50% of the kinase activity was blocked (IC₅₀) was calculated for various cyclin/CDK pairs. The results are indicated in Table I.

TABLE I

Inhibition of cyclin dependent kinase complexes by p27-p16 fusion protein

inhibitor	CDK4/cycl in D1	CDK2/cycl in E	CDK2/cycl in A	cdc2/cyclin B
p27-p16	25 nm	30 nm	25 nm	15 nm
p27	63 nm	52 nm	65 nm	20 nm
p16	250 nm	>500 nm	>500 nm	>500 nm

nm = nanomolar

[0217] Moreover, the inhibition constant, K_i for the inhibition of CDK4/cyclin D1 by p27-p16 fusion protein was determined to be 23 nm, compared to a K_i of 75 nm for p16 inhibition of the same CDK4 complex.

[0218] Other exemplary fusion proteins were derived as follows:

[0219] (i) a "p16(GS)p27" fusion protein was generated to include, N to C terminal, the entire coding sequence of p16, fused in frame with a (Gly,Ser)₃ linker and then the full coding sequence of p27. The nucleotide sequence for the fusion gene encoding the p16(GS)p27

protein is provided in SEQ ID No. 4, with the corresponding amino acid sequence being designated by SEQ ID No. 5; and

[0220] (ii) a "p16p27" fusion protein was generated to include, N to C terminal, the entire coding sequence of p16, fused in frame the full coding sequence of p27 (no (Gly₄Ser)₃ linker). The nucleotide sequence for the fusion gene encoding the p16p27 protein is provided in SEQ ID No. 6, with the corresponding amino acid sequence being designated by SEQ ID No. 7.

[0221] In other embodiments, we have constructed fusion inhibitors that contain smaller portions of p27. In these p27 derivatives we have deleted sequences that may play a role in p27 degradation. The idea here that we may increase in vivo efficacy by increasing potency and stability, e.g., by rendering the fusion resistant to ubiquitin-mediated degradation. The truncated p27 protein (12aa-177aa) is shown in SEQ ID No. 16, which provides a polypeptide of the formula EcoR1-ATG-HA epitope-p27(12aa-177aa)-Stop-Not1. Still another truncated variant of p27 is provided in SEQ ID No. 18, and encodes a polypeptide of the formula EcoR1-ATG-HA epitope-p27(25aa-97aa)-STOP-Not1.

[0222] Utilizing these p27 fragments, we have constructed p27/p16 fusion proteins of the formula EcoR1-ATG-HA epitope-p27(12aa-177aa)-p16(2aa-155aa)-STOP-Not1, which is shown in SEQ ID No. 20, and of the formula EcoR1-ATG-HA epitope-p27(12aa-177aa)-(Gly₄Ser)₃ hinge-p16((2aa-155aa)-STOP-Not1, shown in SEQ ID No. 22.

[0223] Still other p27/p16 fusion with the truncated p27 is EcoR1-ATG-HA epitope-p27(25aa-93aa)-p16(2aa-155aa)-STOP-Not1, shown in SEQ ID No. 24, and EcoR1-ATG-HA epitope-p27(25aa-93aa)-(Gly₄Ser)₃ hinge-p16(2aa-155aa)-STOP-Not1, shown in SEQ ID No. 26.

[0224] All of the above-cited references and publications are hereby incorporated by reference.

Equivalents

[0225] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 34

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1420 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: both
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 4..1176

(ix) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 4..24
- (D) OTHER INFORMATION: /product= "poly-His_Tag"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAT ATG CAC CAC CAT CAC CAT CAC TCA AAC GTG CGA GTG TCT AAC GGG	48
Met His His His His His Ser Asn Val Arg Val Ser Asn Gly	
1 5 10 15	
AGC CCT AGC CTG GAG CGG ATG GAC GCC AGG CAG GCG GAG CAC CCC AAG	96
Ser Pro Ser Leu Glu Arg Met Asp Ala Arg Gln Ala Glu His Pro Lys	
20 25 30	
CCC TCG GCC TGC AGG AAC CTC TTC GGC CCG GTG GAC CAC GAA GAG TTA	144
Pro Ser Ala Cys Arg Asn Leu Phe Gly Pro Val Asp His Glu Glu Leu	
35 40 45	
ACC CGG GAC TTG GAG AAG CAC TGC AGA GAC ATG GAA GAG GCG AGC CAG	192
Thr Arg Asp Leu Glu Lys His Cys Arg Asp Met Glu Glu Ala Ser Gln	
50 55 60	
CGC AAG TGG AAT TTC GAT TTT CAG AAT CAC AAA CCC CTA GAG GGC AAG	240

-continued

Arg	Lys	Trp	Asn	Phe	Asp	Phe	Gln	Asn	His	Lys	Pro	Leu	Glu	Gly	Lys	
65						70					75					
TAC	GAG	TGG	CAA	GAG	GTG	GAG	AAG	GGC	AGC	TTG	CCC	GAG	TTC	TAC	TAC	288
Tyr	Glu	Trp	Gln	Glu	Val	Glu	Lys	Gly	Ser	Leu	Pro	Glu	Phe	Tyr	Tyr	
80					85					90					95	
AGA	CCC	CCG	CGG	CCC	CCC	AAA	GGT	GCC	TGC	AAG	GTG	CCG	GCG	CAG	GAG	336
Arg	Pro	Pro	Arg	Pro	Pro	Lys	Gly	Ala	Cys	Lys	Val	Pro	Ala	Gln	Glu	
				100					105					110		
AGC	CAG	GAT	GTC	AGC	GGG	AGC	CGC	CCG	GCG	GCG	CCT	TTA	ATT	GGG	GCT	384
Ser	Gln	Asp	Val	Ser	Gly	Ser	Arg	Pro	Ala	Ala	Pro	Leu	Ile	Gly	Ala	
			115					120					125			
CCG	GCT	AAC	TCT	GAG	GAC	ACG	CAT	TTG	GTG	GAC	CCA	AAG	ACT	GAT	CCG	432
Pro	Ala	Asn	Ser	Glu	Asp	Thr	His	Leu	Val	Asp	Pro	Lys	Thr	Asp	Pro	
		130					135					140				
TCG	GAC	AGC	CAG	ACG	GGG	TTA	GCG	GAG	CAA	TGC	GCA	GGA	ATA	AGG	AAG	480
Ser	Asp	Ser	Gln	Thr	Gly	Leu	Ala	Glu	Gln	Cys	Ala	Gly	Ile	Arg	Lys	
	145					150					155					
CGA	CCT	GCA	ACC	GAC	GAT	TCT	TCT	ACT	CAA	AAC	AAA	AGA	GCC	AAC	AGA	528
Arg	Pro	Ala	Thr	Asp	Asp	Ser	Ser	Thr	Gln	Asn	Lys	Arg	Ala	Asn	Arg	
160					165					170					175	
ACA	GAA	GAA	AAT	GTT	TCA	GAC	GGT	TCC	CCA	AAT	GCC	GGT	TCT	GTG	GAG	576
Thr	Glu	Glu	Asn	Val	Ser	Asp	Gly	Ser	Pro	Asn	Ala	Gly	Ser	Val	Glu	
			180					185						190		
CAG	ACG	CCC	AAG	AAG	CCT	GGC	CTC	AGA	AGA	CGT	CAA	ACG	GGT	GGC	GGG	624
Gln	Thr	Pro	Lys	Lys	Pro	Gly	Leu	Arg	Arg	Arg	Gln	Thr	Gly	Gly	Gly	
			195				200						205			
GGC	AGC	GGG	GGT	GGC	GGT	TCC	GGC	GGG	GGT	GGA	TCC	GAA	TTC	TGC	GGC	672
Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Phe	Cys	Gly	
		210				215					220					
CGC	GCG	TGC	GCT	CGG	CGG	CTG	CGG	AGA	GGG	GAG	AGC	ATG	CAG	CGG	GCG	720
Arg	Ala	Cys	Ala	Arg	Arg	Leu	Arg	Arg	Gly	Glu	Ser	Met	Gln	Arg	Ala	
	225					230					235					
GCG	GGG	AGC	AGC	ATG	GAG	CCT	TCG	GCT	GAC	TGG	CTG	GCC	ACG	GCC	GCG	768
Ala	Gly	Ser	Ser	Met	Glu	Pro	Ser	Ala	Asp	Trp	Leu	Ala	Thr	Ala	Ala	
240				245						250				255		
GCC	CGG	GGT	CGG	GTA	GAG	GAG	GTG	CGG	GCG	CTG	CTG	GAG	GCG	GTG	GCG	816
Ala	Arg	Gly	Arg	Val	Glu	Glu	Val	Arg	Ala	Leu	Leu	Glu	Ala	Val	Ala	
			260					265					270			
CTG	CCC	AAC	GCA	CCG	AAT	AGT	TAC	GGT	CGG	AGG	CCG	ATC	CAG	GTC	ATG	864
Leu	Pro	Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg	Arg	Pro	Ile	Gln	Val	Met	
		275					280						285			
ATG	ATG	GGC	AGC	GCC	CGA	GTG	GCG	GAG	CTG	CTG	CTG	CTC	CAC	GGC	GCG	912
Met	Met	Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu	Leu	Leu	Leu	His	Gly	Ala	
		290				295						300				
GAG	CCC	AAC	TGC	GCC	GAC	CCC	GCC	ACT	CTC	ACC	CGA	CCC	GTG	CAC	GAC	960
Glu	Pro	Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu	Thr	Arg	Pro	Val	His	Asp	
	305					310					315					
GCT	GCC	CGG	GAG	GGC	TTC	CTG	GAC	ACG	CTG	GTG	GTG	CTG	CAC	CGG	GCC	1008
Ala	Ala	Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu	Val	Val	Leu	His	Arg	Ala	
320				325						330				335		
GGG	GCG	CGG	CTG	GAC	GTG	CGC	GAT	GCC	TGG	GGC	CGT	CTG	CCC	GTG	GAC	1056
Gly	Ala	Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp	Gly	Arg	Leu	Pro	Val	Asp	
			340					345					350			
CTG	GCT	GAG	GAG	CTG	GGC	CAT	CGC	GAT	GTC	GCA	CGG	TAC	CTG	CGC	GCG	1104
Leu	Ala	Glu	Glu	Leu	Gly	His	Arg	Asp	Val	Ala	Arg	Tyr	Leu	Arg	Ala	
		355				360						365				
GCT	GCG	GGG	GGC	ACC	AGA	GGC	AGT	AAC	CAT	GCC	GCG	ATA	GAT	GCC	GCG	1152

-continued

Ala	Ala	Gly	Gly	Thr	Arg	Gly	Ser	Asn	His	Ala	Arg	Ile	Asp	Ala	Ala	
		370					375					380				
GAA	GGT	CCC	TCA	GAC	ATC	CCC	GAT	TGAAAGAACC	AGAGAGGCTC	TGAGAAACCT						1206
Glu	Gly	Pro	Ser	Asp	Ile	Pro	Asp									
		385					390									
CGGGAAACTT	AGATCATCAG	TCACCGAAGG	TCCTACAGGG	CCACAACCTGC	CCCCGCCACA											1266
ACCCACCCCG	CTTTCGTAGT	TTTCATTAG	AAAATAGAGC	TTTAAAAAT	GTCTGCCTT											1326
TTAACGTAGA	TATAAGCCTT	CCCCCACTAC	CGTAAATGTC	CATTATATC	ATTTTITATA											1386
TATTCTTATA	AAAATGTAAA	AAAGAAAACT	CGAG													1420

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 391 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	His	His	His	His	His	His	Ser	Asn	Val	Arg	Val	Ser	Asn	Gly	Ser	
1					5				10					15		
Pro	Ser	Leu	Glu	Arg	Met	Asp	Ala	Arg	Gln	Ala	Glu	His	Pro	Lys	Pro	
		20					25						30			
Ser	Ala	Cys	Arg	Asn	Leu	Phe	Gly	Pro	Val	Asp	His	Glu	Glu	Leu	Thr	
		35					40					45				
Arg	Asp	Leu	Glu	Lys	His	Cys	Arg	Asp	Met	Glu	Glu	Ala	Ser	Gln	Arg	
		50				55				60						
Lys	Trp	Asn	Phe	Asp	Phe	Gln	Asn	His	Lys	Pro	Leu	Glu	Gly	Lys	Tyr	
		65			70				75					80		
Glu	Trp	Gln	Glu	Val	Glu	Lys	Gly	Ser	Leu	Pro	Glu	Phe	Tyr	Tyr	Arg	
		85							90					95		
Pro	Pro	Arg	Pro	Pro	Lys	Gly	Ala	Cys	Lys	Val	Pro	Ala	Gln	Glu	Ser	
		100					105					110				
Gln	Asp	Val	Ser	Gly	Ser	Arg	Pro	Ala	Ala	Pro	Leu	Ile	Gly	Ala	Pro	
		115				120					125					
Ala	Asn	Ser	Glu	Asp	Thr	His	Leu	Val	Asp	Pro	Lys	Thr	Asp	Pro	Ser	
		130				135					140					
Asp	Ser	Gln	Thr	Gly	Leu	Ala	Glu	Gln	Cys	Ala	Gly	Ile	Arg	Lys	Arg	
		145			150				155					160		
Pro	Ala	Thr	Asp	Asp	Ser	Ser	Thr	Gln	Asn	Lys	Arg	Ala	Asn	Arg	Thr	
		165					170						175			
Glu	Glu	Asn	Val	Ser	Asp	Gly	Ser	Pro	Asn	Ala	Gly	Ser	Val	Glu	Gln	
		180					185						190			
Thr	Pro	Lys	Lys	Pro	Gly	Leu	Arg	Arg	Arg	Gln	Thr	Gly	Gly	Gly	Gly	
		195					200				205					
Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Phe	Cys	Gly	Arg	
		210				215					220					
Ala	Cys	Ala	Arg	Arg	Leu	Arg	Arg	Gly	Glu	Ser	Met	Gln	Arg	Ala	Ala	
		225			230				235					240		
Gly	Ser	Ser	Met	Glu	Pro	Ser	Ala	Asp	Trp	Leu	Ala	Thr	Ala	Ala	Ala	
			245					250					255			
Arg	Gly	Arg	Val	Glu	Glu	Val	Arg	Ala	Leu	Leu	Glu	Ala	Val	Ala	Leu	

-continued

260	265	270
Pro Asn Ala Pro Asn Ser Tyr Gly Arg Arg Pro Ile Gln Val Met Met		
275	280	285
Met Gly Ser Ala Arg Val Ala Glu Leu Leu Leu Leu His Gly Ala Glu		
290	295	300
Pro Asn Cys Ala Asp Pro Ala Thr Leu Thr Arg Pro Val His Asp Ala		
305	310	315
Ala Arg Glu Gly Phe Leu Asp Thr Leu Val Val Leu His Arg Ala Gly		
325	330	335
Ala Arg Leu Asp Val Arg Asp Ala Trp Gly Arg Leu Pro Val Asp Leu		
340	345	350
Ala Glu Glu Leu Gly His Arg Asp Val Ala Arg Tyr Leu Arg Ala Ala		
355	360	365
Ala Gly Gly Thr Arg Gly Ser Asn His Ala Arg Ile Asp Ala Ala Glu		
370	375	380
Gly Pro Ser Asp Ile Pro Asp		
385	390	

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 51 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GCGGCCGCGTC ATATGCACCA CCATCACCAT CACTCAAACG TCGAGTGTCT T 51

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 96 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GCGGCCGCGCG TCGACTCGGC CGAATTCGGA TCCACCCCGC CCGGAACCGC CACCCCGCT 60

GCCCCCGCCA CCCGTTTGAC GTCTTCTGAG GCCAGG 96

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1143 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..1140

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG GGA TAC CCT TAT GAT GTG CCA GAT TAT GCC GAT CCG GCG GCG GGG 48

-continued

Met	Gly	Tyr	Pro	Tyr	Asp	Val	Pro	Asp	Tyr	Ala	Asp	Pro	Ala	Ala	Gly	
1				5					10						15	
AGC	AGC	ATG	GAG	CCT	TCG	GCT	GAC	TGG	CTG	GCC	ACG	GCC	GCG	GCC	CGG	96
Ser	Ser	Met	Glu	Pro	Ser	Ala	Asp	Trp	Leu	Ala	Thr	Ala	Ala	Ala	Arg	
			20					25					30			
GGT	CGG	GTA	GAG	GAG	GTG	CGG	GCG	CTG	CTG	GAG	GCG	GGG	GCG	CTG	CCC	144
Gly	Arg	Val	Glu	Glu	Val	Arg	Ala	Leu	Leu	Glu	Ala	Gly	Ala	Leu	Pro	
			35					40					45			
AAC	GCA	CCG	AAT	AGT	TAC	GGT	CGG	AGG	CCG	ATC	CAG	GTC	ATG	ATG	ATG	192
Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg	Arg	Pro	Ile	Gln	Val	Met	Met	Met	
			50					55				60				
GGC	AGC	GCC	CGA	GTG	GCG	GAG	CTG	CTG	CTG	CTC	CAC	GCG	GCG	GAG	CCC	240
Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu	Leu	Leu	His	Gly	Ala	Glu	Pro		
						70				75				80		
AAC	TGC	GCC	GAC	CCC	GCC	ACT	CTC	ACC	CGA	CCC	GTG	CAC	GAC	GCT	GCC	288
Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu	Thr	Arg	Pro	Val	His	Asp	Ala	Ala	
						85				90				95		
CGG	GAG	GGC	TTC	CTG	GAC	ACG	CTG	GTG	GTG	CTG	CAC	CGG	GCC	GGG	GCG	336
Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu	Val	Val	Leu	His	Arg	Ala	Gly	Ala	
				100				105						110		
CGG	CTG	GAC	GTG	GCG	GAT	GCC	TGG	GGC	CGT	CTG	CCC	GTG	GAC	CTG	GCT	384
Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp	Gly	Arg	Leu	Pro	Val	Asp	Leu	Ala	
				115				120						125		
GAG	GAG	CTG	GGC	CAT	GCG	GAT	GTC	GCA	CGG	TAC	CTG	GCG	GCG	GCT	GCG	432
Glu	Glu	Leu	Gly	His	Arg	Asp	Val	Ala	Arg	Tyr	Leu	Arg	Ala	Ala	Ala	
			130					135				140				
GGG	GGC	ACC	AGA	GGC	AGT	AAC	CAT	GCC	CGC	ATA	GAT	GCC	GCG	GAA	GGT	480
Gly	Gly	Thr	Arg	Gly	Ser	Asn	His	Ala	Arg	Ile	Asp	Ala	Ala	Glu	Gly	
						150				155				160		
CCC	TCA	GAC	ATC	CCC	GAT	GGT	GGC	GGG	GGC	AGC	GGG	GGT	GGC	GGT	TCC	528
Pro	Ser	Asp	Ile	Pro	Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	
						165				170				175		
GGC	GGG	GGT	GGA	TCC	GTC	GAG	TCA	AAC	GTG	CGA	GTG	TCT	AAC	GGG	GCG	576
Gly	Gly	Gly	Gly	Ser	Val	Glu	Ser	Asn	Val	Arg	Val	Ser	Asn	Gly	Arg	
						180				185				190		
CCT	AGC	CTG	GAG	CGG	ATG	GAC	GCC	AGG	CAG	GCG	GAG	CAC	CCC	AAG	CCC	624
Pro	Ser	Leu	Glu	Arg	Met	Asp	Ala	Arg	Gln	Ala	Glu	His	Pro	Lys	Pro	
								200					205			
TCG	GCC	TGC	AGG	AAC	CTC	TTC	GGC	CCG	GTG	GAC	CAC	GAA	GAG	TTA	ACC	672
Ser	Ala	Cys	Arg	Asn	Leu	Phe	Gly	Pro	Val	Asp	His	Glu	Glu	Leu	Thr	
								215						220		
CGG	GAC	TTG	GAG	AAG	CAC	TGC	AGA	GAC	ATG	GAA	GAG	GCG	AGC	CAG	GCG	720
Arg	Asp	Leu	Glu	Lys	His	Cys	Arg	Asp	Met	Glu	Glu	Ala	Ser	Gln	Arg	
						230				235				240		
AAG	TGG	AAT	TTC	GAT	TTT	CAG	AAT	CAC	AAA	CCC	CTA	GAG	GGC	AAG	TAC	768
Lys	Trp	Asn	Phe	Asp	Phe	Gln	Asn	His	Lys	Pro	Leu	Glu	Gly	Lys	Tyr	
						245				250				255		
GAG	TGG	CAA	GAG	GTG	GAG	AAG	GGC	AGC	TTG	CCC	GAG	TTC	TAC	TAC	AGA	816
Glu	Trp	Gln	Glu	Val	Glu	Lys	Gly	Ser	Leu	Pro	Glu	Phe	Tyr	Tyr	Arg	
						260				265				270		
CCC	CCG	CGG	CCC	CCC	AAA	GGT	GCC	TGC	AAG	GTG	CCG	GCG	CAG	GAG	AGC	864
Pro	Pro	Arg	Pro	Pro	Lys	Gly	Ala	Cys	Lys	Val	Pro	Ala	Gln	Glu	Ser	
						275				280				285		
CAG	GAT	GTC	AGC	GGG	AGC	CGC	CCG	GCG	GCG	CCT	TTA	ATT	GGG	GCT	CCG	912
Gln	Asp	Val	Ser	Gly	Ser	Arg	Pro	Ala	Ala	Pro	Leu	Ile	Gly	Ala	Pro	
						295					300					
GCT	AAC	TCT	GAG	GAC	ACG	CAT	TTG	GTG	GAC	CCA	AAG	ACT	GAT	CCG	TCG	960

-continued

Ala	Asn	Ser	Glu	Asp	Thr	His	Leu	Val	Asp	Pro	Lys	Thr	Asp	Pro	Ser	
305					310					315					320	
GAC	AGC	CAG	ACG	GGG	TTA	CGC	GAG	CAA	TGC	GCA	GGA	ATA	AGG	AAG	CGA	1008
Asp	Ser	Gln	Thr	Gly	Leu	Ala	Glu	Gln	Cys	Ala	Gly	Ile	Arg	Lys	Arg	
				325					330					335		
CCT	GCA	ACC	GAC	GAT	TCT	TCT	ACT	CAA	AAC	AAA	AGA	GCC	AAC	AGA	ACA	1056
Pro	Ala	Thr	Asp	Asp	Ser	Ser	Thr	Gln	Asn	Lys	Arg	Ala	Asn	Arg	Thr	
				340				345					350			
GAA	GAA	AAT	GTT	TCA	GAC	GGT	TCC	CCA	AAT	GCC	GGT	TCT	GTG	GAG	CAG	1104
Glu	Glu	Asn	Val	Ser	Asp	Gly	Ser	Pro	Asn	Ala	Gly	Ser	Val	Glu	Gln	
		355					360					365				
ACG	CCC	AAG	AAG	CCT	GGC	CTC	AGA	AGA	CGT	CAA	ACG	TAA				1143
Thr	Pro	Lys	Lys	Pro	Gly	Leu	Arg	Arg	Arg	Gln	Thr					
	370					375					380					

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 380 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Gly	Tyr	Pro	Tyr	Asp	Val	Pro	Asp	Tyr	Ala	Asp	Pro	Ala	Ala	Gly
1				5					10					15	
Ser	Ser	Met	Glu	Pro	Ser	Ala	Asp	Trp	Leu	Ala	Thr	Ala	Ala	Ala	Arg
		20						25					30		
Gly	Arg	Val	Glu	Glu	Val	Arg	Ala	Leu	Leu	Glu	Ala	Gly	Ala	Leu	Pro
		35					40					45			
Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg	Arg	Pro	Ile	Gln	Val	Met	Met	Met
	50					55				60					
Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu	Leu	Leu	Leu	His	Gly	Ala	Glu	Pro
	65			70					75				80		
Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu	Thr	Arg	Pro	Val	His	Asp	Ala	Ala
		85						90					95		
Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu	Val	Val	Leu	His	Arg	Ala	Gly	Ala
	100					105						110			
Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp	Gly	Arg	Leu	Pro	Val	Asp	Leu	Ala
	115					120						125			
Glu	Glu	Leu	Gly	His	Arg	Asp	Val	Ala	Arg	Tyr	Leu	Arg	Ala	Ala	Ala
	130					135					140				
Gly	Gly	Thr	Arg	Gly	Ser	Asn	His	Ala	Arg	Ile	Asp	Ala	Ala	Glu	Gly
	145				150					155				160	
Pro	Ser	Asp	Ile	Pro	Asp	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
		165					170						175		
Gly	Gly	Gly	Gly	Ser	Val	Glu	Ser	Asn	Val	Arg	Val	Ser	Asn	Gly	Arg
		180						185					190		
Pro	Ser	Leu	Glu	Arg	Met	Asp	Ala	Arg	Gln	Ala	Glu	His	Pro	Lys	Pro
	195					200					205				
Ser	Ala	Cys	Arg	Asn	Leu	Phe	Gly	Pro	Val	Asp	His	Glu	Glu	Leu	Thr
	210					215					220				
Arg	Asp	Leu	Glu	Lys	His	Cys	Arg	Asp	Met	Glu	Glu	Ala	Ser	Gln	Arg
	225			230						235				240	

-continued

Lys	Trp	Asn	Phe	Asp	Phe	Gln	Asn	His	Lys	Pro	Leu	Glu	Gly	Lys	Tyr	
				245					250					255		
Glu	Trp	Gln	Glu	Val	Glu	Lys	Gly	Ser	Leu	Pro	Glu	Phe	Tyr	Tyr	Arg	
			260					265					270			
Pro	Pro	Arg	Pro	Pro	Lys	Gly	Ala	Cys	Lys	Val	Pro	Ala	Gln	Glu	Ser	
		275					280					285				
Gln	Asp	Val	Ser	Gly	Ser	Arg	Pro	Ala	Ala	Pro	Leu	Ile	Gly	Ala	Pro	
	290					295					300					
Ala	Asn	Ser	Glu	Asp	Thr	His	Leu	Val	Asp	Pro	Lys	Thr	Asp	Pro	Ser	
305					310					315					320	
Asp	Ser	Gln	Thr	Gly	Leu	Ala	Glu	Gln	Cys	Ala	Gly	Ile	Arg	Lys	Arg	
			325						330					335		
Pro	Ala	Thr	Asp	Asp	Ser	Ser	Thr	Gln	Asn	Lys	Arg	Ala	Asn	Arg	Thr	
			340					345					350			
Glu	Glu	Asn	Val	Ser	Asp	Gly	Ser	Pro	Asn	Ala	Gly	Ser	Val	Glu	Gln	
		355					360					365				
Thr	Pro	Lys	Lys	Pro	Gly	Leu	Arg	Arg	Arg	Gln	Thr					
	370					375					380					

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1098 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..1095

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG	GGA	TAC	CCT	TAT	GAT	GTG	CCA	GAT	TAT	GCC	GAT	CCG	GCG	GCG	GGG	48
Met	Gly	Tyr	Pro	Tyr	Asp	Val	Pro	Asp	Tyr	Ala	Asp	Pro	Ala	Ala	Gly	
1				5					10					15		
AGC	AGC	ATG	GAG	CCT	TCG	GCT	GAC	TGG	CTG	GCC	ACG	GCC	GCG	GCC	CGG	96
Ser	Ser	Met	Glu	Pro	Ser	Ala	Asp	Trp	Leu	Ala	Thr	Ala	Ala	Ala	Arg	
			20					25					30			
GGT	CGG	GTA	GAG	GAG	GTG	CGG	GCG	CTG	CTG	GAG	GCG	GGG	GCG	CTG	CCC	144
Gly	Arg	Val	Glu	Glu	Val	Arg	Ala	Leu	Leu	Glu	Ala	Gly	Ala	Leu	Pro	
		35				40					45					
AAC	GCA	CCG	AAT	AGT	TAC	GGT	CGG	AGG	CCG	ATC	CAG	GTC	ATG	ATG	ATG	192
Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg	Arg	Pro	Ile	Gln	Val	Met	Met	Met	
	50					55				60						
GGC	AGC	GCC	CGA	GTG	GCG	GAG	CTG	CTG	CTG	CTC	CAC	GCG	GCG	GAG	CCC	240
Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu	Leu	Leu	Leu	His	Gly	Ala	Glu	Pro	
	65			70					75					80		
AAC	TGC	GCC	GAC	CCC	GCC	ACT	CTC	ACC	CGA	CCC	GTG	CAC	GAC	GCT	GCC	288
Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu	Thr	Arg	Pro	Val	His	Asp	Ala	Ala	
			85						90					95		
CGG	GAG	GCG	TTC	CTG	GAC	ACG	CTG	GTG	GTG	CTG	CAC	CGG	GCC	GGG	GCG	336
Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu	Val	Val	Leu	His	Arg	Ala	Gly	Ala	
			100				105						110			
CGG	CTG	GAC	GTG	GCG	GAT	GCC	TGG	GGC	CGT	CTG	CCC	GTG	GAC	CTG	GCT	384
Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp	Gly	Arg	Leu	Pro	Val	Asp	Leu	Ala	
	115					120						125				

-continued

GAG Glu 130	GAG Glu 130	CTG Leu 130	GGC Gly 130	CAT His 130	CGC Arg 135	GAT Asp 135	GTC Val 135	GCA Ala 135	CGG Arg 135	TAC Tyr 140	CTG Leu 140	CGC Arg 140	GCT Ala 140	CGC Ala 140	432	
GGG Gly 145	GGC Gly 145	ACC Thr 145	AGA Arg 145	GGC Gly 150	AGT Ser 150	AAC Asn 150	CAT His 150	GCC Ala 155	CGC Arg 155	ATA Ile 155	GAT Asp 155	GCC Ala 160	GCG Ala 160	GAA Glu 160	GGT Gly 160	480
CCC Pro 165	TCA Ser 165	GAC Asp 165	ATC Ile 165	CCC Pro 165	GAT Asp 170	GTC Val 170	GAG Glu 170	TCA Ser 170	AAC Asn 170	GTG Val 175	CGA Arg 175	GTG Val 175	TCT Ser 175	AAC Asn 175	GGG Gly 175	528
CGC Arg 180	CCT Pro 180	AGC Ser 180	CTG Leu 180	GAG Glu 180	CGG Arg 185	ATG Met 185	GAC Asp 185	GCC Ala 185	AGG Arg 185	CAG Gln 190	GCG Ala 190	GAG Glu 190	CAC His 190	CCC Pro 190	AAG Lys 190	576
CCC Pro 195	TCG Ser 195	GCC Ala 195	TGC Cys 195	AGG Arg 200	AAC Asn 200	CTC Leu 200	TTC Phe 200	GGC Gly 200	CCG Pro 205	GTG Val 205	GAC Asp 205	CAC His 205	GAA Glu 205	GAG Glu 205	TTA Leu 205	624
ACC Thr 210	CGG Arg 210	GAC Asp 210	TTG Leu 210	GAG Glu 215	AAG Lys 215	CAC His 215	TGC Cys 215	AGA Arg 220	GAC Asp 220	ATG Met 220	GAA Glu 220	GAG Glu 220	GCG Ala 220	AGC Ser 220	CAG Gln 220	672
CGC Arg 225	AAG Lys 225	TGG Trp 225	AAT Asn 230	TTC Phe 230	GAT Asp 230	TTT Phe 230	CAG Gln 235	AAT Asn 235	CAC His 235	AAA Lys 235	CCC Pro 240	CTA Leu 240	GAG Glu 240	GGC Gly 240	ARG Lys 240	720
TAC Tyr 245	GAG Glu 245	TGG Trp 245	CAA Gln 245	GAG Val 250	GTG Glu 250	GAG Lys 250	AAG Gly 250	GGC Ser 250	AGC Leu 250	TTG Pro 255	CCC Glu 255	GAG Phe 255	TTC Tyr 255	TAC Tyr 255	TAC Tyr 255	768
AGA Arg 260	CCC Pro 260	CCG Pro 260	CGG Arg 260	CCC Pro 265	CCC Pro 265	AAA Lys 265	GGT Gly 265	GCC Ala 265	TGC Cys 265	AAG Lys 270	GTG Val 270	CCG Pro 270	GCG Ala 270	CAG Gln 270	GAG Glu 270	816
AGC Ser 275	CAG Gln 275	GAT Val 275	GTC Ser 275	AGG Gly 280	GGG Ser 280	AGC Ser 280	CGC Arg 280	CCG Pro 285	GCG Ala 285	GCG Ala 285	CCT Pro 285	TTA Leu 285	ATT Ile 285	GGG Gly 285	GCT Ala 285	864
CCG Pro 290	GCT Ala 290	AAC Asn 295	TCT Ser 295	GAG Glu 295	GAC Asp 295	ACG Thr 295	CAT His 300	TTG Leu 300	GTG Val 300	GAC Asp 300	CCA Pro 300	AAG Lys 300	ACT Thr 300	GAT Asp 300	CCG Pro 300	912
TCG Ser 305	GAC Asp 305	AGC Ser 310	CAG Gln 310	ACG Thr 310	GGG Gly 310	TTA Leu 310	GCG Ala 315	GAG Glu 315	CAA Gln 315	TGC Cys 315	GCA Ala 315	GGA Gly 320	ATA Ile 320	AGG Arg 320	AAG Lys 320	960
CGA Arg 325	CCT Pro 325	GCA Ala 325	ACC Thr 325	GAC Asp 330	GAT Ser 330	TCT Ser 330	TCT Ser 330	ACT Thr 330	CAA Gln 330	AAC Asn 335	AAA Lys 335	AGA Arg 335	GCC Ala 335	AAC Asn 335	AGA Arg 335	1008
ACA Thr 340	GAA Glu 340	GAA Glu 340	AAT Asn 340	GTT Val 345	TCA Ser 345	GAC Asp 345	GGT Gly 345	TCC Ser 345	CCA Pro 350	AAT Asn 350	GCC Ala 350	GGT Gly 350	TCT Ser 350	GTG Val 350	GAG Glu 350	1056
CAG Gln 355	ACG Thr 355	CCC Pro 355	AAG Lys 355	AAG Lys 360	CCT Pro 360	GGC Gly 360	CTC Leu 360	AGA Arg 365	AGA Arg 365	CGT Arg 365	CAA Gln 365	ACG Thr 365	TAA Thr 365			1098

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 365 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Asp Pro Ala Ala Gly
 1 5 10 15
 Ser Ser Met Glu Pro Ser Ala Asp Trp Leu Ala Thr Ala Ala Ala Arg

-continued

20					25					30					
Gly	Arg	Val	Glu	Glu	Val	Arg	Ala	Leu	Leu	Glu	Ala	Gly	Ala	Leu	Pro
		35					40					45			
Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg	Arg	Pro	Ile	Gln	Val	Met	Met	Met
		50					55					60			
Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu	Leu	Leu	Leu	His	Gly	Ala	Glu	Pro
		65					70					75			80
Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu	Thr	Arg	Pro	Val	His	Asp	Ala	Ala
			85						90					95	
Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu	Val	Val	Leu	His	Arg	Ala	Gly	Ala
			100					105					110		
Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp	Gly	Arg	Leu	Pro	Val	Asp	Leu	Ala
		115					120					125			
Glu	Glu	Leu	Gly	His	Arg	Asp	Val	Ala	Arg	Tyr	Leu	Arg	Ala	Ala	Ala
		130					135					140			
Gly	Gly	Thr	Arg	Gly	Ser	Asn	His	Ala	Arg	Ile	Asp	Ala	Ala	Glu	Gly
		145					150					155			160
Pro	Ser	Asp	Ile	Pro	Asp	Val	Glu	Ser	Asn	Val	Arg	Val	Ser	Asn	Gly
			165						170					175	
Arg	Pro	Ser	Leu	Glu	Arg	Met	Asp	Ala	Arg	Gln	Ala	Glu	His	Pro	Lys
			180					185					190		
Pro	Ser	Ala	Cys	Arg	Asn	Leu	Phe	Gly	Pro	Val	Asp	His	Glu	Glu	Leu
		195					200					205			
Thr	Arg	Asp	Leu	Glu	Lys	His	Cys	Arg	Asp	Met	Glu	Glu	Ala	Ser	Gln
		210					215					220			
Arg	Lys	Trp	Asn	Phe	Asp	Phe	Gln	Asn	His	Lys	Pro	Leu	Glu	Gly	Lys
		225					230					235			240
Tyr	Glu	Trp	Gln	Glu	Val	Glu	Lys	Gly	Ser	Leu	Pro	Glu	Phe	Tyr	Tyr
			245					250						255	
Arg	Pro	Pro	Arg	Pro	Pro	Lys	Gly	Ala	Cys	Lys	Val	Pro	Ala	Gln	Glu
			260				265					270			
Ser	Gln	Asp	Val	Ser	Gly	Ser	Arg	Pro	Ala	Ala	Pro	Leu	Ile	Gly	Ala
		275					280					285			
Pro	Ala	Asn	Ser	Glu	Asp	Thr	His	Leu	Val	Asp	Pro	Lys	Thr	Asp	Pro
		290					295					300			
Ser	Asp	Ser	Gln	Thr	Gly	Leu	Ala	Glu	Gln	Cys	Ala	Gly	Ile	Arg	Lys
		305					310					315			320
Arg	Pro	Ala	Thr	Asp	Asp	Ser	Ser	Thr	Gln	Asn	Lys	Arg	Ala	Asn	Arg
			325					330					335		
Thr	Glu	Glu	Asn	Val	Ser	Asp	Gly	Ser	Pro	Asn	Ala	Gly	Ser	Val	Glu
			340					345					350		
Gln	Thr	Pro	Lys	Lys	Pro	Gly	Leu	Arg	Arg	Arg	Gln	Thr			
		355					360					365			

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 75 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CATATGGGTG GCTGCCGTGG CGATATGTTC GGTTCGGGTG CTCCTCCAAA AAAGAAGAGA 60
 AAGGTAGCTG GATTG 75

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Gly Gly Cys Arg Gly Asp Met Phe Gly Cys Gly Ala Pro Pro Lys
 1 5 10 15
 Lys Lys Arg Lys Val Ala Gly Phe
 20

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 225 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CATATGGAGC CAGTAGATCC TAGACTAGAG CCCTGGAAGC ATCCAGGAAG TCAGCCTAAA 60
 ACTGCTTGTA CCAATTGCTA TTGTA AAAAG TGTTCCTTTC ATTGCCAAGT TTGTTTCATA 120
 ACAAAGCCC TTGGCATCTC CTATGGCAGG AAGAAGCGGA GACAGCGACG AAGACCTCCT 180
 CAAGGCAGTC AGACTCATCA AGTTTCTCTA AGTAAGCAAG GATTG 225

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 72 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Glu Pro Val Asp Pro Arg Leu Glu Pro Trp Lys His Pro Gly Ser
 1 5 10 15
 Gln Pro Lys Thr Ala Cys Thr Asn Cys Tyr Cys Lys Lys Cys Cys Phe
 20 25 30
 His Cys Gln Val Cys Phe Ile Thr Lys Ala Leu Gly Ile Ser Tyr Gly
 35 40 45
 Arg Lys Lys Arg Arg Gln Arg Arg Pro Pro Gln Gly Ser Gln Thr
 50 55 60
 His Gln Val Ser Leu Ser Lys Gln
 65 70

(2) INFORMATION FOR SEQ ID NO:13:

-continued

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 912 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

CATATGACCT CTCGCCGCTC CGTGAAGTCG GGTCCGCGGG AGGTTCCGCG CGATGAGTAC   60
GAGGATCTGT ACTACACCCC GTCTTCAGGT ATGGCGAGTC CCGATAGTCC GCCTGACACC   120
TCCCGCCGTG GCGCCCTACA GACACGCTCG CGCCAGAGGG GCGAGGTCCG TTTCGTCCAG   180
TACGACGAGT CGGATTATGC CCTCTACGGG GGCTCGTCAT CCGAAGACGA CGAACACCCG   240
GAGGTCCCCC GGACGCGGCG TCCCGTTTCC GGGGCGGTTT TGTCCGGCCC GGGGCCTGCG   300
CGGGCGCCTC CGCCACCCGC TGGGTCCGGA GGGGCCGGAC GCACACCCAC CACCGCCCCC   360
CGGGCCCCCC GAACCCAGCG GGTGGCGACT AAGGCCCCCG CGGCCCCGGC GCGCGAGACC   420
ACCCGCGGCA GGAATCCGGC CCAGCCAGAA TCCGCCGCAC TCCAGACGC CCCCGCGTCG   480
ACGGCGCCAA CCCGATCCAA GACACCCGCG CAGGGGCTGG CCAGAAAGCT GCACCTTIAGC   540
ACCGCCCCCC CAAACCCCGA CGCGCCATGG ACCCCCCGGG TGGCCGGCTT TAACAAGCGC   600
GTCTTCTGCG CCGCGGTCGG GCGCCTGGCG GCCATGCATG CCCGGATGGC GCGGCTCCAG   660
CTCTGGGACA TGTCGCGTCC GCGCACAGAC GAAGACCTCA ACGAACTCCT TGGCATCACC   720
ACCATCCGCG TGACGGTCTG CGAGGGCAAA AACCTGCTTC AGCGCGCCAA CGAGTTGGTG   780
AATCCAGACG TGGTGCAGGA CGTCGACGCG GCCACGGCGA CTCGAGGGCG TTCTGCGGCG   840
TCGCGCCCCA CCGAGCGACC TCGAGCCCCA GCCCGCTCCG CTTCTCGCCC CAGACGGCCC   900
GTCGAGGAAT TC                                          912

```

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 301 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

Met Thr Ser Arg Arg Ser Val Lys Ser Gly Pro Arg Glu Val Pro Arg
1      5      10      15
Asp Glu Tyr Glu Asp Leu Tyr Tyr Thr Pro Ser Ser Gly Met Ala Ser
20     25     30
Pro Asp Ser Pro Pro Asp Thr Ser Arg Arg Gly Ala Leu Gln Thr Arg
35     40     45
Ser Arg Gln Arg Gly Glu Val Arg Phe Val Gln Tyr Asp Glu Ser Asp
50     55     60
Tyr Ala Leu Tyr Gly Ser Ser Ser Glu Asp Asp Glu His Pro Glu
65     70     75     80
Val Pro Arg Thr Arg Arg Pro Val Ser Gly Ala Val Leu Ser Gly Pro
85     90     95
Gly Pro Ala Arg Ala Pro Pro Pro Pro Ala Gly Ser Gly Gly Ala Gly
100    105    110

```

-continued

Arg Thr Pro Thr Thr Ala Pro Arg Ala Pro Arg Thr Gly Arg Val Ala
 115 120 125
 Thr Lys Ala Pro Ala Ala Pro Ala Ala Glu Thr Thr Arg Gly Arg Lys
 130 135 140
 Ser Ala Gln Pro Glu Ser Ala Ala Leu Pro Asp Ala Pro Ala Ser Thr
 145 150 155 160
 Ala Pro Thr Arg Ser Lys Thr Pro Ala Gln Gly Leu Ala Arg Lys Leu
 165 170 175
 His Phe Ser Thr Ala Pro Pro Asn Pro Asp Ala Pro Trp Thr Pro Arg
 180 185 190
 Val Ala Gly Phe Asn Lys Arg Val Phe Cys Ala Ala Val Gly Arg Leu
 195 200 205
 Ala Ala Met His Ala Arg Met Ala Ala Val Gln Leu Trp Asp Met Ser
 210 215 220
 Arg Pro Arg Thr Asp Glu Asp Leu Asn Glu Leu Leu Gly Ile Thr Thr
 225 230 235 240
 Ile Arg Val Thr Val Cys Glu Gly Lys Asn Leu Leu Gln Arg Ala Asn
 245 250 255
 Glu Leu Val Asn Pro Asp Val Val Gln Asp Val Asp Ala Ala Thr Ala
 260 265 270
 Thr Arg Gly Arg Ser Ala Ala Ser Arg Pro Thr Glu Arg Pro Arg Ala
 275 280 285
 Pro Ala Arg Ser Ala Ser Arg Pro Arg Arg Pro Val Glu
 290 295 300

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 120 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CATATGGACG TCGACGCGGC CACGGCGACT CGAGGGCGTT CTGCGGCGTC GCGCCCCACC 60
 GAGCGACCTC GAGCCCCAGC CCGCTCCGCT TCTCGCCCCA GACGGCCCGT CGAGGAATTC 120

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Met Asp Val Asp Ala Ala Thr Ala Thr Arg Gly Arg Ser Ala Ala Ser
 1 5 10 15
 Arg Pro Thr Glu Arg Pro Arg Ala Pro Ala Arg Ser Ala Ser Arg Pro
 20 25 30
 Arg Arg Pro Val Glu
 35

-continued

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 557 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

```

GAATTCGCCG CCACCATGGG ATACCCTTAT GATGTGCCAG ATTATGCCAG CCTGGAGCGG      60
ATGGACGCCA GGCAGGCGGA GCACCCCAAG CCCTCGGCCT GCAGGAACCT CTCGGGCCCG      120
GTGGACCACG AAGAGTTAAC CCGGACTTG GAGAAGCACT GCAGAGACAT GGAAGAGGCG      180
AGCCAGCGCA AGTGAATTTC CGATTTTCAG AATCACAAC CCCTAGAGGG CAAGTACGAG      240
TGGCAAGAGG TGGAGAAGGG CAGCTTGCCC GAGTTCTACT ACAGACCCCC GCGGCCCCCC      300
AAAGTGCCCT GCAAGGTGCC GGCGCAGGAG AGCCAGGATG TCAGCGGGAG CCGCCCGGCG      360
GCGCCTTTAA TTGGGGCTCC GGCTAACTCT GAGGACACGC ATTTGGTGGA CCCAAAGACT      420
GATCCGTCGG ACAGCCAGAC GGGGTTAGCG GAGCAATGCG CAGGAATAAG GAAGCGACCT      480
GCAACCGACG ATTCTTCTAC TCAAACAAA AGAGCCAACA GAACAGAAGA AATGTTTCA      540
GACGGTTAGG CGGCCGCG                                     557

```

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 167 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

```

Met Ser Leu Glu Arg Met Asp Ala Arg Gln Ala Glu His Pro Lys Pro
1      5      10      15
Ser Ala Cys Arg Asn Leu Phe Gly Pro Val Asp His Glu Glu Leu Thr
20     25     30
Arg Asp Leu Glu Lys His Cys Arg Asp Met Glu Glu Ala Ser Gln Arg
35     40     45
Lys Trp Asn Phe Asp Phe Gln Asn His Lys Pro Leu Glu Gly Lys Tyr
50     55     60
Glu Trp Gln Glu Val Glu Lys Gly Ser Leu Pro Glu Phe Tyr Tyr Arg
65     70     75     80
Pro Pro Arg Pro Pro Lys Gly Ala Cys Lys Val Pro Ala Gln Glu Ser
85     90     95
Gln Asp Val Ser Gly Ser Arg Pro Ala Ala Pro Leu Ile Gly Ala Pro
100    105    110
Ala Asn Ser Glu Asp Thr His Leu Val Asp Pro Lys Thr Asp Pro Ser
115    120    125
Asp Ser Gln Thr Gly Leu Ala Glu Gln Cys Ala Gly Ile Arg Lys Arg
130    135    140
Pro Ala Thr Asp Asp Ser Ser Thr Gln Asn Lys Arg Ala Asn Arg Thr
145    150    155    160

```

-continued

Glu Glu Asn Val Ser Asp Gly
165

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 266 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```
GAATTCGCCG CCACCATGGG ATACCCTTAT GATGTGCCAG ATTATGCCAA GCCCTCGGCC      60
TGCAGGAACC TCTTCGGCCC GGTGGACCAC GAAGAGTTAA CCCGGGACTT GGAGAAGCAC      120
TGCAGAGACA TGAAGAGGCC GAGCCAGCGC AAGTGAATT TCGATTTTCA GAATCACAAA      180
CCCTAGAGG GCAAGTACGA GTGGCAAGAG GTGGAGAAGG GCAGCTTGCC CGAGTTCTAC      240
TACAGACCCC CGCGGTAGGC GGCCGC                                           266
```

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 70 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

```
Met Lys Pro Ser Ala Cys Arg Asn Leu Phe Gly Pro Val Asp His Glu
1           5           10          15
Glu Leu Thr Arg Asp Leu Glu Lys His Cys Arg Asp Met Glu Glu Ala
20          25          30
Ser Gln Arg Lys Trp Asn Phe Asp Phe Gln Asn His Lys Pro Leu Glu
35          40          45
Gly Lys Tyr Glu Trp Gln Glu Val Glu Lys Gly Ser Leu Pro Glu Phe
50          55          60
Tyr Tyr Arg Pro Pro Arg
65          70
```

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1028 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```
GAATTCGCCG CCACCATGGG ATACCCTTAT GATGTGCCAG ATTATGCCAG CCTGGAGCGG      60
ATGGACGCCA GGCAGGCGGA GCACCCCAAG CCCTCGGCCT GCAGGAACCT CTTGGGCCCG      120
GTGGACCACG AAGAGTTAAC CCGGACTTGA GAGAAGCACT GCAGAGACAT GGAAGAGGCG      180
AGCCAGCGCA AGTGAATTTT CGATTTTCAG AATCACAAC CCCTAGAGGG CAAGTACGAG      240
TGGCAAGAGG TGGAGAAGGG CAGCTTGCCC GAGTTCTACT ACAGACCCCC GCGGCCCCCC      300
```

-continued

```

AAAGGTGCCT GCAAGGTGCC GGCGCAGGAG AGCCAGGATG TCAGCGGGAG CCGCCCGGCG 360
GCGCCTTTAA TTGGGGCTCC GGCTAACTCT GAGGACACGC ATTTGGTGGA CCCAAAGACT 420
GATCCGTCGG ACAGCCAGAC GGGGTTAGCG GAGCAATGCG CAGGAATAAG GAAGCGACCT 480
GCAACCGACG ATTCTTCTAC TCAAAACAAA AGAGCCAACA GAACAGAAGA AAATGTTTCA 540
GACGGTGTGCG AGGATCCGGC GGCGGGGAGC AGCATGGAGC CTTCGGGTGA CTGGCTGGCC 600
ACGGCCGCGG CCCGGGGTCG GGTAGAGGAG GTGCGGGCGC TGCTGGAGGC GGGGGCGCTG 660
CCCAACGCAC CGAATAGTTA CGGTCGGAGG CCGATCCAGG TCATGATGAT GGGCAGCGCC 720
CGAGTGGCGG AGCTGCTGCT GCTCCACGGC GCGGAGCCCA ACTGCGCCGA CCCCGCCACT 780
CTCACCCGAC CCGTGACAGA CGCTGCCCGG GAGGGCTTCC TGGACACGCT GGTGGTGCTG 840
CACCGGGCCG GGGCGCGGCT GGACGTGCGC GATGCCTGGG GCCGTCTGCC CGTGGACCTG 900
GCTGAGGAGC TGGGCCATCG CGATGTCGCA CGGTACCTGC GCGCGGCTGC GGGGGGCACC 960
AGAGGCAGTA ACCATGCCCG CATAGATGCC GCGGAAGGTC CCTCAGACAT CCCCGATTGA 1020
GCGGCGCGC 1028

```

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 334 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser Leu Glu Arg Met
1           5           10           15
Asp Ala Arg Gln Ala Glu His Pro Lys Pro Ser Ala Cys Arg Asn Leu
20          25          30
Phe Gly Pro Val Asp His Glu Glu Leu Thr Arg Asp Leu Glu Lys His
35          40          45
Cys Arg Asp Met Glu Glu Ala Ser Gln Arg Lys Trp Asn Phe Asp Phe
50          55          60
Gln Asn His Lys Pro Leu Glu Gly Lys Tyr Glu Trp Gln Glu Val Glu
65          70          75          80
Lys Gly Ser Leu Pro Glu Phe Tyr Tyr Arg Pro Pro Arg Pro Pro Lys
85          90          95
Gly Ala Cys Lys Val Pro Ala Gln Glu Ser Gln Asp Val Ser Gly Ser
100         105         110
Arg Pro Ala Ala Pro Leu Ile Gly Ala Pro Ala Asn Ser Glu Asp Thr
115         120         125
His Leu Val Asp Pro Lys Thr Asp Pro Ser Asp Ser Gln Thr Gly Leu
130         135         140
Ala Glu Gln Cys Ala Gly Ile Arg Lys Arg Pro Ala Thr Asp Asp Ser
145         150         155         160
Ser Thr Gln Asn Lys Arg Ala Asn Arg Thr Glu Glu Asn Val Ser Asp
165         170         175
Gly Val Glu Asp Pro Ala Ala Gly Ser Ser Met Glu Pro Ser Ala Asp
180         185         190

```


-continued

Trp	Leu	Ala	Thr	Ala	Ala	Ala	Arg	Gly	Arg	Val	Glu	Glu	Val	Arg	Ala
	195						200					205			
Leu	Leu	Glu	Ala	Gly	Ala	Leu	Pro	Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg
	210					215					220				
Arg	Pro	Ile	Gln	Val	Met	Met	Met	Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu
	225				230					235					240
Leu	Leu	Leu	His	Gly	Ala	Glu	Pro	Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu
			245						250					255	
Thr	Arg	Pro	Val	His	Asp	Ala	Ala	Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu
			260					265					270		
Val	Val	Leu	His	Arg	Ala	Gly	Ala	Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp
		275				280						285			
Gly	Arg	Leu	Pro	Val	Asp	Leu	Ala	Glu	Glu	Leu	Gly	His	Arg	Asp	Val
	290					295					300				
Ala	Arg	Tyr	Leu	Arg	Ala	Ala	Ala	Gly	Gly	Thr	Arg	Gly	Ser	Asn	His
	305				310					315					320
Ala	Arg	Ile	Asp	Ala	Ala	Glu	Gly	Pro	Ser	Asp	Ile	Pro	Asp		
			325						330						

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1073 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GAATTCGCCG	CCACCATGGG	ATACCCCTTAT	GATGTGCCAG	ATTATGCCAG	CCTGGAGCGG	60
ATGGACGCCA	GGCAGGCGGA	GCACCCCAAG	CCCTCGGCCT	GCAGGAACCT	CTTCGGCCCCG	120
GTGGACCACG	AAGAGTTAAC	CCGGGACTTG	GAGAAGCACT	GCAGAGACAT	GGAAGAGGCG	180
AGCCAGCGCA	AGTGAATTT	CGATTTTCAG	AATCACAAC	CCCTAGAGGG	CAAGTACGAG	240
TGGCAAGAGG	TGGAGAAGGG	CAGCTTGCCC	GAGTTCTACT	ACAGACCCCC	GCGGCCCCCC	300
AAAGGTGCCT	GCAAGGTGCC	GGCGCAGGAG	AGCCAGGATG	TCAGCGGGAG	CCGCCCGGCG	360
GCGCCTTTAA	TTGGGGCTCC	GGCTAACTCT	GAGGACACGC	ATTTGGTGGA	CCCAAAGACT	420
GATCCGTCGG	ACAGCCAGAG	GGGGTTAGCG	GAGCAATGCG	CAGGAATAAG	GAAGCGACCT	480
GCAACCGACG	ATTCTTCTAC	TCAAAACAAA	AGAGCCAACA	GAACAGAAGA	AAATGTTTCA	540
GACGGTGGTG	GCGGGGGCAG	CGGGGGTGCG	GGTTCCGGCG	GGGGTGGATC	CGTCGAGGAT	600
CCGGCGGCGG	GGAGCAGCAT	GGAGCCTTCG	GCTGACTGGC	TGGCCACGGC	CGCGGCCCGG	660
GGTCGGSTAG	AGGAGGTGCG	GGCGCTGCTG	GAGGCGGGGG	CGCTGCCCAA	CGACCCGAAT	720
AGTTACGGTC	GGAGGCCGAT	CCAGGTCATG	ATGATGGGCA	GCGCCCGAGT	GGCGGAGCTG	780
CTGCTGCTCC	ACGGCGCGGA	GCCCAACTGC	GCCGACCCCG	CCACTCTCAC	CCGACCCGTG	840
CACGACGCTG	CCCGGGAGGG	CTTCCTGGAC	ACGCTGGTGG	TGCTGCACCG	GCCCGGGGCG	900
CGGCTGGACG	TGCGCGATGC	CTGGGGCCGT	CTGCCCSTGG	ACCTGGCTGA	GGAGCTGGGC	960
CATCGCGATG	TCGCACGGTA	CCTGCGCGCG	GCTGCGGGGG	GCACCAGAGG	CAGTAACCAT	1020
GCCCCGATAG	ATGCCGCGGA	AGGTCCTCTA	GACATCCCCG	ATTGAGCGGC	CGC	1073

-continued

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 348 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser Leu Glu Arg Met
 1           5           10           15

Asp Ala Arg Gln Ala Glu His Pro Lys Pro Ser Ala Cys Arg Asn Leu
 20           25           30

Phe Gly Pro Val Asp His Glu Glu Leu Thr Arg Asp Leu Glu Lys His
 35           40           45

Cys Arg Asp Met Glu Glu Ala Ser Gln Arg Lys Trp Asn Phe Asp Phe
 50           55           60

Gln Asn His Lys Pro Leu Glu Gly Lys Tyr Glu Trp Gln Glu Val Glu
 65           70           75           80

Lys Gly Ser Leu Pro Glu Phe Tyr Tyr Arg Pro Pro Arg Pro Pro Lys
 85           90           95

Gly Ala Cys Lys Val Pro Ala Gln Glu Ser Gln Asp Val Ser Gly Ser
100           105           110

Arg Pro Ala Ala Pro Leu Ile Gly Ala Pro Ala Asn Ser Glu Asp Thr
115           120           125

His Leu Val Asp Pro Lys Thr Asp Pro Ser Asp Ser Gln Thr Gly Leu
130           135           140

Ala Glu Gln Cys Ala Gly Ile Arg Lys Arg Pro Ala Thr Asp Asp Ser
145           150           155           160

Ser Thr Gln Asn Lys Arg Ala Asn Arg Thr Glu Glu Asn Val Ser Asp
165           170           175

Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
180           185           190

Val Glu Asp Pro Ala Ala Gly Ser Ser Met Glu Pro Ser Ala Asp Trp
195           200           205

Leu Ala Thr Ala Ala Ala Arg Gly Arg Val Glu Glu Val Arg Ala Leu
210           215           220

Leu Glu Ala Gly Ala Leu Pro Asn Ala Pro Asn Ser Tyr Gly Arg Arg
225           230           235           240

Pro Ile Gln Val Met Met Met Gly Ser Ala Arg Val Ala Glu Leu Leu
245           250           255

Leu Leu His Gly Ala Glu Pro Asn Cys Ala Asp Pro Ala Thr Leu Thr
260           265           270

Arg Pro Val His Asp Ala Ala Arg Glu Gly Phe Leu Asp Thr Leu Val
275           280           285

Val Leu His Arg Ala Gly Ala Arg Leu Asp Val Arg Asp Ala Trp Gly
290           295           300

Leu Pro Val Asp Leu Ala Glu Glu Leu Gly His Arg Asp Val Ala Arg
305           310           315           320

Tyr Leu Arg Ala Ala Ala Gly Gly Thr Arg Gly Ser Asn His Ala Arg
325           330           335

```

-continued

Ile Asp Ala Ala Glu Gly Pro Ser Asp Ile Pro Asp
340 345

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 737 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

```

GAATTCGCCG CCACCATGGG ATACCCTTAT GATGTGCCAG ATTATGCCAA GCCCTCGGCC      60
TGCAGGAACC TCTTCGGCCC GGTGGACCAC GAAGACTTAA CCCGGGACTT GGAGAAGCAC      120
TGCAGAGACA TGAAGAGGCG GAGCCAGCGC AAGTGAATT TCGATTTTCA GAATCACAAA      180
CCCTTAGAGG GCAAGTACGA GTGGCAAGAG GTGGAGAAGG GCAGCTTGCC CGAGTTCTAC      240
TACAGACCCC CGCGGGTCGA GGATCCGGCG GCGGGGAGCA GCATGGAGCC TTCGGCTGAC      300
TGGCTGGCCA CGGCCGCGCG CCGGGGTCGG GTAGAGGAGG TCGGGGCGCT GCTGGAGGCG      360
GGGGCGCTGC CCAACGCACC GAATAGTTAC GGTCCGAGGC CGATCCAGGT CATGATGATG      420
GGCAGCGCCC GAGTGGCGGA GCTGCTGCTG CTCACGGCG CGGAGCCCAA CTGCGCCGAC      480
CCCGCCACTC TCACCCGACC CGTGACGAC GCTGCCCGGG AGGGCTTCCT GGACACGCTG      540
GTGGTGCTGC ACCGGGCGCG GGCGCGGCTG GACGTGCGCG ATGCCTGGGG CCGTCTGCCC      600
GTGGACCTGG CTGAGGAGCT GGGCCATCGC GATGTGCGAC GGTACCTGCG CGCGGGTGCG      660
GGGGGCACCA GAGGCAGTAA CCATGCCCGC ATAGATGCCG CGGAAGGTCC CTCAGACATC      720
CCCGATTGAG CGGCCCGC

```

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 237 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

```

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Lys Pro Ser Ala Cys
1      5      10      15
Arg Asn Leu Phe Gly Pro Val Asp His Glu Glu Leu Thr Arg Asp Leu
20     25     30
Glu Lys His Cys Arg Asp Met Glu Glu Ala Ser Gln Arg Lys Trp Asn
35     40     45
Phe Asp Phe Gln Asn His Lys Pro Leu Glu Gly Lys Tyr Glu Trp Gln
50     55     60
Glu Val Glu Lys Gly Ser Leu Pro Glu Phe Tyr Tyr Arg Pro Pro Arg
65     70     75     80
Val Glu Asp Pro Ala Ala Gly Ser Ser Met Glu Pro Ser Ala Asp Trp
85     90     95
Leu Ala Thr Ala Ala Ala Arg Gly Arg Val Glu Glu Val Arg Ala Leu
100    105    110

```

-continued

Leu	Glu	Ala	Gly	Ala	Leu	Pro	Asn	Ala	Pro	Asn	Ser	Tyr	Gly	Arg	Arg
	115						120						125		
Pro	Ile	Gln	Val	Met	Met	Met	Gly	Ser	Ala	Arg	Val	Ala	Glu	Leu	Leu
	130						135					140			
Leu	Leu	His	Gly	Ala	Glu	Pro	Asn	Cys	Ala	Asp	Pro	Ala	Thr	Leu	Thr
	145				150					155					160
Arg	Pro	Val	His	Asp	Ala	Ala	Arg	Glu	Gly	Phe	Leu	Asp	Thr	Leu	Val
				165					170					175	
Val	Leu	His	Arg	Ala	Gly	Ala	Arg	Leu	Asp	Val	Arg	Asp	Ala	Trp	Gly
			180					185					190		
Arg	Leu	Pro	Val	Asp	Leu	Ala	Glu	Glu	Leu	Gly	His	Arg	Asp	Val	Ala
		195					200					205			
Arg	Tyr	Leu	Arg	Ala	Ala	Gly	Gly	Thr	Arg	Gly	Ser	Asn	His	Ala	
	210					215					220				
Arg	Ile	Asp	Ala	Ala	Glu	Gly	Pro	Ser	Asp	Ile	Pro	Asp			
	225				230					235					

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 782 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GAATTCGCCG CCACCATGGG ATACCCTTAT GATGTGCCAG ATTATGCCAA GCCCTCGGCC	60
TGCAGGAACC TCTTCGGCCC GGTGGACCAC GAAGAGTTAA CCCGGGACTT GGAGAAGCAC	120
TGCAGAGACA TGGAAGAGGC GAGCCAGCGC AAGTGAATT TCGATTTTCA GAATCACAAA	180
CCCCTAGAGG GCAASTACGA GTGGCAAGAG GTGGAGAAGG GCAGCTTGCC CGAGTTCTAC	240
TACAGACCCC CGCGGGGTGG CGGGGCGAGC GGGGGTGGCG GTTCCGGCGG GGGTGGATCC	300
GTCGAGGATC CGGCGGCGGG GAGCAGCATG GAGCCTTCGG CTGACTGGCT GGCCACGGCC	360
GCGGCCCGGG GTCGGGTAGA GGAGGTGCGG GCGCTGCTGG AGGCGGGGGG GCTGCCCAAC	420
GCACCGAATA GTTACGGTCG GAGGCCGATC CAGGTCATGA TGATGGGCAG CGCCCGAGTG	480
GCGGAGCTGC TGCTGCTCCA CGGCGCGGAG CCCAACTGCG CCGACCCCGC CACTCTCACC	540
CGACCCGTGC ACGACGCTGC CCGGAGGGC TTCCTGGACA CGCTGGTGGT GCTGCACCGG	600
GCCGGGGCGC GGCTGGACGT GCGCGATGCC TGGGGCCGTC TGCCCGTGGA CCTGGCTGAG	660
GAGCTGGGCC ATCGCGATGT CGCACGGTAC CTGCGCGCGG CTGCGGGGGG CACCAGAGGC	720
AGTAACCATG CCCGCATAGA TGCCGCGGAA GGTCCCTCAG ACATCCCCGA TTGAGCGGCC	780
GC	782

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 252 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

```

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Lys Pro Ser Ala Cys
1           5           10           15
Arg Asn Leu Phe Gly Pro Val Asp His Glu Glu Leu Thr Arg Asp Leu
20           25           30
Glu Lys His Cys Arg Asp Met Glu Glu Ala Ser Gln Arg Lys Trp Asn
35           40           45
Phe Asp Phe Gln Asn His Lys Pro Leu Glu Gly Lys Tyr Glu Trp Gln
50           55           60
Glu Val Glu Lys Gly Ser Leu Pro Glu Phe Tyr Tyr Arg Pro Pro Arg
65           70           75           80
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Val
85           90           95
Glu Asp Pro Ala Ala Gly Ser Ser Met Glu Pro Ser Ala Asp Trp Leu
100          105          110
Ala Thr Ala Ala Ala Arg Gly Arg Val Glu Glu Val Arg Ala Leu Leu
115          120          125
Glu Ala Gly Ala Leu Pro Asn Ala Pro Asn Ser Tyr Gly Arg Arg Pro
130          135          140
Ile Gln Val Met Met Met Gly Ser Ala Arg Val Ala Glu Leu Leu Leu
145          150          155          160
Leu His Gly Ala Glu Pro Asn Cys Ala Asp Pro Ala Thr Leu Thr Arg
165          170          175
Pro Val His Asp Ala Ala Arg Glu Gly Phe Leu Asp Thr Leu Val Val
180          185          190
Leu His Arg Ala Gly Ala Arg Leu Asp Val Arg Asp Ala Trp Gly Arg
195          200          205
Leu Pro Val Asp Leu Ala Glu Glu Leu Gly His Arg Asp Val Ala Arg
210          215          220
Tyr Leu Arg Ala Ala Ala Gly Gly Thr Arg Gly Ser Asn His Ala Arg
225          230          235          240
Ile Asp Ala Ala Glu Gly Pro Ser Asp Ile Pro Asp
245          250

```

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

```

Cys Phe Ile Thr Lys Ala Leu Gly Ile Ser Tyr Gly Arg Lys Lys Arg
1           5           10           15
Arg Gln Arg Arg Arg Pro Pro Gln Gly Ser
20           25

```

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 12 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Cys Met His Ile Glu Ser Leu Asp Ser Tyr Thr Cys
1 5 10

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Cys Met Tyr Ile Glu Ala Leu Asp Lys Tyr Ala Cys
1 5 10

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Xaa Xaa Xaa Glu Ala Ala Leu Ala Glu Ala Leu Ala Glu Ala Leu Ala
1 5 10 15

Glu Ala Leu Ala Glu Ala Leu Ala Glu Ala Leu Glu Ala Leu Ala Ala
20 25 30

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Gly Asn Ala Ala Ala Arg Arg
1 5

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Cys Asp Pro Gly Tyr Ile Gly Ser Arg Cys
1 5 10

We claim:

1. A recombinant transfection system, comprising
 - (i) a first gene construct comprising a coding sequence encoding an inhibitory polypeptide comprising at least one CDK-binding motif for binding and inhibiting activation of a cyclin dependent kinase (cdk), which coding sequence is operably linked to a transcriptional regulatory sequence for causing expression of the first polypeptide in eukaryotic cells,
 - (ii) a second gene construct comprising a coding sequence encoding an endothelialization polypeptide which promotes endothelialization, and
 - (iii) a gene delivery composition for delivering the gene constructs to a cell and causing the cell to be transfected with the gene construct.
2. The transfection system of claim 1, wherein the CDK-binding motif is a CDK-binding motif of a CDK inhibitor protein.
3. The transfection system of claim 2, wherein the CDK inhibitor protein is an INK4 protein.
4. The transfection system of claim 3, wherein the INK4 protein is selected from the group consisting of p15, p16, p18 and p19.
5. The transfection system of claim 2, wherein the CDK inhibitor protein is a CIP protein.
6. The transfection system of claim 5, wherein the CIP protein is selected from the group consisting of p21^{CIP1}, p27^{KIP1}, and p57^{KIP2}.
7. The transfection system of claim 1, wherein the CDK-binding motif comprises tandemly arranged ankyrin-like sequences.
8. The transfection system of claim 1, wherein the CDK-binding motif comprises a p21/p27 inhibitory domain.
9. The transfection system of claim 1, wherein the inhibitory polypeptide is a fusion protein comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases.
10. The transfection system of claim 1, wherein the endothelialization polypeptide stimulates endothelial cell proliferation.
11. The transfection system of claim 1, wherein the endothelialization polypeptide stimulates migration of endothelial cells to a wound site.
12. The transfection system of claim 1, wherein the endothelialization polypeptide is selected from the group consisting of angiogenic basic fibroblast growth factors (bFGF), acid fibroblast growth factor (aFGF), vascular endothelial growth factor (VEGF), platelet derived growth factor (PDGF), vascular permeability growth factor (VPF), and transforming growth factor beta (TGF-β).
13. The transfection system of claim 1, wherein the first and second gene constructs are provided as part of a single vector.
14. The transfection system of claim 1, wherein the first and second gene constructs are provided as part of polycistronic message.
15. The transfection system of claim 1, wherein the first and second gene constructs are provided on separate vectors.
16. The recombinant transfection system of claim 1, wherein the gene construct comprises a viral vector.
17. The recombinant transfection system of claim 16, wherein the viral vector is an adenoviral vector.
18. The recombinant transfection system of claim 16, wherein the viral vector is an adeno-associated viral vector.
19. The recombinant transfection system of claim 16, wherein the viral vector is a retroviral vector.
20. The recombinant transfection system of claim 1, wherein the gene delivery composition comprises a recombinant viral particle.
21. The recombinant transfection system of claim 1, wherein the gene delivery composition is selected from the group consisting of a liposome and a poly-cationic nucleic acid binding agent.
22. The recombinant transfection system of claim 1, wherein the gene delivery composition further comprises a pharmaceutically acceptable carrier for administration to an animal.
23. A method for treating an animal having a vascular wound characterized a breach of endothelial integrity and by excessive smooth muscle proliferation, comprising administering the recombinant transfection system of claim 1 to the area of the vascular wound.
24. The method of claim 23, wherein the vascular wound is restenosis.
25. The method of claim 23, wherein the recombinant transfection system is administered by catheter.
26. A nucleic acid encoding a fusion protein comprising a therapeutic polypeptide sequence from an intracellular protein which alter a biological process of a cell upon intracellular localization of the fusion protein, and a transcellular polypeptide sequence for promoting transecytosis of the fusion protein across a cell surface membrane and into a cell.
27. The transfection system of claim 26, wherein the therapeutic polypeptide sequence includes a CDK-binding motif.
28. The transfection system of claim 27, wherein the CDK-binding motif is a CDK-binding motif of a CDK inhibitor protein.
29. The transfection system of claim 28, wherein the CDK inhibitor protein is an INK4 protein.
30. The transfection system of claim 29, wherein the INK4 protein is selected from the group consisting of p15, p16, p18 and p19.
31. The transfection system of claim 28, wherein the CDK inhibitor protein is a CIP protein.
32. The transfection system of claim 31, wherein the CIP protein is selected from the group consisting of p21^{CIP1}, p27^{KIP1}, and p57^{KIP2}.
33. The transfection system of claim 27, wherein the CDK-binding motif comprises tandemly arranged ankyrin-like sequences.
34. The transfection system of claim 27, wherein the CDK-binding motif comprises a p21/p27 inhibitory domain.
35. The transfection system of claim 27, wherein the inhibitory polypeptide is a fusion protein comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases.
36. The transfection system of claim 26, wherein the therapeutic polypeptide sequence is from a tumor suppressor, a transcription factor, a signal transduction protein, an antiviral protein or a metal chelating protein.
37. The transfection system of claim 36, wherein the therapeutic polypeptide sequence includes a polypeptide sequence of a tumor suppressor selected from the group consisting of p53, Rb and Rb-like proteins.

38. The transfection system of claim 36, wherein the therapeutic polypeptide sequence includes a polypeptide sequence of a signal transduction protein selected from the group consisting of tubby, a DOT protein, a Bcl protein, and an IκB protein.

39. The transfection system of claim 38, wherein the Bcl protein is Bcl-2 or Bcl-x.

40. The transfection system of claim 26, wherein the biological process is selected from the group consisting of proliferation, differentiation, cell death, gene expression, protein stability, calcium mobilization, ion permeability, phosphorylation of intracellular proteins, metabolism of inositol phosphates, and metabolism of nucleosides.

41. The transfection system of claim 26, wherein the fusion protein inhibits protein-protein interactions or protein-nucleic acid interactions between proteins and nucleic acids endogenous to the cell.

42. The transfection system of claim 26, wherein the transcellular polypeptide sequence is an internalizing peptide.

43. The transfection system of claim 42, wherein the internalizing peptide is derived from a polypeptide selected from the group consisting of antepennepedia protein, HIV transactivating (TAT) protein, mastoparan, melittin, bombolittin, delta hemolysin, pardaxin, Pseudomonas exotoxin A, clathrin, Diphtheria toxin and C9 complement protein.

44. The transfection system of claim 42, wherein the internalizing peptide is derived from antepennepedia protein.

45. The transfection system of claim 42, wherein the internalizing peptide is derived from HIV transactivating (TAT) protein.

46. The transfection system of claim 26, wherein the transcellular polypeptide sequence is an accessory peptide sequence which enhances interaction of the fusion protein with a cell surface membrane.

47. The transfection system of claim 42, wherein the accessory peptide includes an RGD sequence.

48. The recombinant transfection system of claim 26, wherein the gene construct comprises a viral vector.

49. The recombinant transfection system of claim 48, wherein the viral vector is an adenoviral vector.

50. The recombinant transfection system of claim 48, wherein the viral vector is an adeno-associated viral vector.

51. The recombinant transfection system of claim 48, wherein the viral vector is a retroviral vector.

52. The recombinant transfection system of claim 26, wherein the gene delivery composition comprises a recombinant viral particle.

53. The recombinant transfection system of claim 26, wherein the gene delivery composition is selected from the group consisting of a liposome and a poly-cationic nucleic acid binding agent.

54. The recombinant transfection system of claim 26, wherein the gene delivery composition further comprises a pharmaceutically acceptable carrier for administration to an animal.

55. A fusion protein comprising at least one CDK-binding motif for binding and inhibiting activation of a cyclin dependent kinase (cdk), and a transcellular polypeptide sequence for promoting transcytosis of the fusion protein across a cell surface membrane and into a cell.

56. A method for treating an animal for unwanted cell proliferation, comprising administering to the animal the recombinant transfection system of claim 26 or the fusion protein of claim 55.

57. A nucleic acid encoding a fusion protein comprising a first polypeptide sequence having therapeutic activity, and a transcellular polypeptide sequence for promoting transcytosis of the fusion protein across a cell surface membrane and into a cell.

58. A fusion protein encoded by the nucleic acid of claim 57.

59. A nucleic acid comprising a nucleotide sequence encoding a chimeric polypeptide comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases.

60. The nucleic acid of claim 59, which nucleic acid comprises the coding sequence designated in one of SEQ ID Nos. 1, 4, 6, 16, 18, 20, 22, 24 and 26.

61. A recombinant transfection system, comprising

(i) a gene construct including a nucleic acid encoding a chimeric polypeptide comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases, and operably linked to a transcriptional regulatory sequence for causing expression of the chimeric polypeptide in eukaryotic cells, and

(ii) a gene delivery composition for delivering the gene construct to a cell and causing the cell to be transfected with the gene construct.

62. A chimeric polypeptide comprising CDK-binding motifs from two or more different proteins which bind to cyclin dependent kinases.

56. The chimeric polypeptide of claim 55, which chimeric polypeptide comprises the fusion sequence designated in SEQ ID Nos. 2, 5, 7, 17, 19, 21, 23, 25 and 27.

* * * * *